

Renormalization and asymptotic expansion of Dirac's polarized vacuum

Philippe Gravejat ¹, Mathieu Lewin ², Éric Séré ³

Abstract

We perform rigorously the charge renormalization of the so-called reduced Bogoliubov-Dirac-Fock (rBDF) model. This nonlinear theory, based on the Dirac operator, describes atoms and molecules while taking into account vacuum polarization effects. We consider the total physical density ρ_{ph} including both the external density of a nucleus and the self-consistent polarization of the Dirac sea, but no 'real' electron. We show that ρ_{ph} admits an asymptotic expansion to any order in powers of the physical coupling constant α_{ph} , provided that the ultraviolet cut-off behaves as $\Lambda \sim e^{3\pi(1-Z_3)/2\alpha_{\text{ph}}} \gg 1$. The renormalization parameter $0 < Z_3 < 1$ is defined by $Z_3 = \alpha_{\text{ph}}/\alpha$ where α is the bare coupling constant. The coefficients of the expansion of ρ_{ph} are independent of Z_3 , as expected. The first order term gives rise to the well-known Uehling potential, whereas the higher order terms satisfy an explicit recursion relation.

1 Introduction and main result

Renormalization is an essential tool in Quantum Electrodynamics (QED) [9, 2, 21]. The purpose of this paper is to perform rigorously the charge renormalization of a nonlinear approximation of QED, the reduced Bogoliubov-Dirac-Fock (rBDF) theory that was studied before in [15, 16, 19, 18, 17, 13]. This model, based on the Dirac operator, describes atoms and molecules while taking into account vacuum polarization effects. It does not need any mass renormalization, hence it is a theory simple enough for an investigation of charge renormalization in full detail.

Before turning to our specific Dirac model, let us quickly recall the spirit of renormalization. A physical theory usually aims at predicting physical observables in terms of the parameters in the model. Sometimes, interesting quantities are divergent and it is necessary to introduce cut-offs. For electrons the parameters are their mass m and their charge e (or rather the coupling constant $\alpha = e^2$). Predicted physical quantities are then functions $F(m, \alpha, \Lambda)$ where Λ is the regularization parameter. Mass and charge are also physical observables and renormalization occurs when their values predicted by the theory are different from their 'bare' values

$$m_{\text{ph}} = m_{\text{ph}}(m, \alpha, \Lambda) \neq m \text{ and/or } \alpha_{\text{ph}} = \alpha_{\text{ph}}(m, \alpha, \Lambda) \neq \alpha. \quad (1)$$

In this case the parameters m and α are not observable in contrast with $m_{\text{ph}} = m_{\text{ph}}(m, \alpha, \Lambda)$ and $\alpha_{\text{ph}} = \alpha_{\text{ph}}(m, \alpha, \Lambda)$ which have to be set equal to their experimental values. The relation (1) has to be inverted, in order to express the bare parameters in terms of the physical ones

$$m = m(m_{\text{ph}}, \alpha_{\text{ph}}, \Lambda) \text{ and } \alpha = \alpha(m_{\text{ph}}, \alpha_{\text{ph}}, \Lambda). \quad (2)$$

¹Centre de Mathématiques Laurent Schwartz (UMR 7640), École Polytechnique, 91128 Palaiseau Cedex, France. E-mail: gravejat@math.polytechnique.fr

²CNRS & Laboratoire de Mathématiques, UMR 8088, Université de Cergy-Pontoise, 2, avenue Adolphe Chauvin, 95302 Cergy-Pontoise Cedex, FRANCE. E-mail: Mathieu.Lewin@math.cnrs.fr

³Centre de Recherche en Mathématiques de la Décision, Université Paris Dauphine, Place du Maréchal De Lattre De Tassigny, 75775 Paris Cedex 16, France. E-mail: sere@ceremade.dauphine.fr

This allows to express any observable quantity F as a function \tilde{F} of the physical parameters and the cut-off Λ

$$\tilde{F}(m_{\text{ph}}, \alpha_{\text{ph}}, \Lambda) = F(m(m_{\text{ph}}, \alpha_{\text{ph}}, \Lambda), \alpha(m_{\text{ph}}, \alpha_{\text{ph}}, \Lambda), \Lambda). \quad (3)$$

A possible definition of renormalizability is that all such observable quantities have a limit when $\Lambda \rightarrow +\infty$, for fixed m_{ph} and α_{ph} .

Important difficulties can be encountered when trying to complete this program:

- The physical quantities m_{ph} and α_{ph} might be non-explicit functions of α and m . The corresponding formulas can then only be inverted perturbatively to any order (usually in α). This is the case in QED [9, 2, 21]. In the model studied in this paper we have $m_{\text{ph}} = m$ and $\alpha_{\text{ph}} \neq \alpha$, hence only the charge has to be renormalized. Furthermore α_{ph} is an explicit function of m , α and Λ (see (8) later). Renormalizing our model is therefore a much easier task than in full QED.
- Even when the bare parameters are explicit functions of the physical ones, these relations can make it impossible to take the limit $\Lambda \rightarrow +\infty$ while keeping m_{ph} and α_{ph} fixed. As we will explain, in our model $(2/3\pi)\alpha_{\text{ph}} \log \Lambda \leq 1$. To deal with this problem, we let Λ depend on α_{ph} and we investigate the asymptotics in the limit $\alpha_{\text{ph}} \rightarrow 0$.

We now turn to the description of our model. The Bogoliubov-Dirac-Fock theory is the Hartree-Fock approximation of QED when photons are neglected [19, 18]. The associated reduced theory is obtained by further neglecting the so-called exchange term. In both models, the system is described by a Hartree-Fock (quasi-free) state in Fock space, which is completely characterized by its one-body density matrix P (an orthogonal projector for pure states), acting on the one-body space. The state P contains both the ‘real’ electrons of the system (that of an atom for instance) and the ‘virtual’ electrons of the Dirac sea, which all interact with each other self-consistently. Therefore, there are always infinitely many particles and P is infinite-rank.

When the exchange term is neglected, a ground state at zero temperature is (formally) a solution of the following self-consistent equation

$$\begin{cases} P = \chi_{(-\infty, \mu)}(D) + \delta, \\ D = D^0 + \alpha(\rho_{P-\frac{1}{2}} - \nu) * \frac{1}{|x|}. \end{cases} \quad (4)$$

Here $D^0 = \boldsymbol{\alpha} \cdot (-i\nabla) + \beta$ is the free Dirac operator [32] acting on the Hilbert space $\mathfrak{H} := L^2(\mathbb{R}^2, \mathbb{C}^4)$. For the sake of simplicity we have chosen units in which the speed of light is $c = 1$ and, as the model does not need any mass renormalization, we have taken $m = 1$ for the mass of the electrons. The second term in the formula of D is the Coulomb potential induced by both a fixed external density of charge ν (modeling for instance a smeared nucleus) and the self-consistent density $\rho_{P-\frac{1}{2}}$ of the system (see below). In (4), α is the bare coupling constant that will be renormalized later and $\mu \in (-1, 1)$ is a chemical potential which is chosen to fix the desired total charge of the system. We have added in (4) the possibility of having a density matrix $0 \leq \delta \leq \chi_{\{\mu\}}(D)$ at the Fermi level, as is usually done in reduced Hartree-Fock theory [31]. So the operator P is not necessarily a projector but we still use the letter P for convenience. Later we will restrict ourselves to the case of P being an orthogonal projector.

Equation (4) is well-known in the physical literature. A mean-field equation of the same form (including an exchange term and classical terms accounting for the interactions with photons) was derived by Reinhard, Greiner and Arenhövel [26] from the Schwinger-Dyson equations of QED. Chaix and Iracane [4] later gave a variational interpretation of this equation.

Related models appear in relativistic Density Functional Theory, usually with additional empirical exchange-correlation terms (see e.g. [12, Eq. (6.2)] and [11, Eq. (62)]). Dirac [8] already considered the first order term obtained from (4) in an expansion in powers of α .

Let us now explain the exact meaning of $\rho_{P-1/2}$. The charge density of an operator $A : \mathfrak{H} \rightarrow \mathfrak{H}$ with integral kernel $A(x, y)_{\sigma, \sigma'}$ is formally defined as $\rho_A(x) = \sum_{\sigma=1}^4 A(x, x)_{\sigma, \sigma} = \text{tr}_{\mathbb{C}^4}(A(x, x))$. In usual Hartree-Fock theory, the charge density is $\rho_P(x)$. However, as there are infinitely many particles, this does not make sense here. In (4), the subtraction of half the identity is a convenient way to give a meaning to the density, independently of any reference. One has formally

$$\rho_{P-\frac{1}{2}}(x) = \rho_{\frac{P-P^\perp}{2}}(x) = \frac{1}{2} \sum_{i \geq 1} |\varphi_i^-(x)|^2 - |\varphi_i^+(x)|^2,$$

where $(\varphi_i^-)_{i \geq 1}$ is an orthonormal basis of $P\mathfrak{H}$ and $(\varphi_i^+)_{i \geq 1}$ is an orthonormal basis of $(1-P)\mathfrak{H}$. As was explained in [19], subtracting 1/2 to the density matrix P of the Hartree-Fock state makes the model invariant under charge conjugation.

When there is no external field, $\nu \equiv 0$, equation (4) has an obvious solution for any $\mu \in (-1, 1)$, the Hartree-Fock state made of all electrons with negative energy

$$P = P_-^0 := \chi_{(-\infty, 0)}(D^0),$$

in accordance with Dirac's ideas [5, 6, 7]. Indeed $\rho_{P_-^0-1/2} \equiv 0$, as is seen by writing in the Fourier representation

$$\left(P_-^0 - \frac{1}{2}\right)(p) = -\frac{\boldsymbol{\alpha} \cdot p + \beta}{2\sqrt{1+|p|^2}},$$

and since the Dirac matrices are trace-less. This shows the usefulness of the subtraction of half the identity to P , since the free vacuum P_-^0 now has a vanishing density. For a general state P , we can use this to write (formally)

$$\rho_{P-\frac{1}{2}} = \rho_{P-\frac{1}{2}} - \rho_{P_-^0-\frac{1}{2}} = \rho_{P-P_-^0}.$$

When P belongs to a suitable class of perturbations of P_-^0 (for instance when $P - P_-^0$ is locally trace-class), the density $\rho_{P-P_-^0}$ is a well-defined mathematical object. We will give below natural conditions which guarantee that $P - P_-^0$ has a well-defined density in our context.

In the presence of an external field, $\nu \neq 0$, equation (4) has no solution in any 'reasonable' Banach space [16] and it is necessary to introduce an ultraviolet regularization parameter Λ . The simplest method (although probably not optimal regarding regularity issues [13]) is to impose a cut-off at the level of the Hilbert space, that is to replace \mathfrak{H} by

$$\mathfrak{H}_\Lambda := \{f \in L^2(\mathbb{R}^3, \mathbb{C}^4), \text{supp}(\hat{f}) \subset B(0, \Lambda)\},$$

and to solve, instead of (4), the regularized equation in \mathfrak{H}_Λ

$$\begin{cases} P = \chi_{(-\infty, \mu)}(D) + \delta, \\ D = \Pi_\Lambda \left(D^0 + \alpha(\rho_{P-P_-^0} - \nu) * \frac{1}{|x|} \right) \Pi_\Lambda, \end{cases} \quad (5)$$

where Π_Λ is the orthogonal projector onto \mathfrak{H}_Λ in $\mathfrak{H} = L^2(\mathbb{R}^3, \mathbb{C}^4)$.

Existence of solutions to (5) was proved in [16] for $\mu = 0$ and in [13] for $\mu \neq 0$. The precise statement is the following¹.

¹To be more precise, in [13, Theorem 1], only the existence and uniqueness of minimizers of the reduced BDF functional are stated. Elementary arguments based on convexity allow to deduce Theorem 1 from the results of [13].

Theorem 1 (Existence of self-consistent solutions to (5) [16, 13]). *Assume that $\alpha \geq 0$, $\Lambda > 0$ and $\mu \in (-1, 1)$ are given. Let ν be in the so-called Coulomb space*

$$\mathcal{C} := \left\{ f : \mathbb{R}^3 \rightarrow \mathbb{R}, \text{ s.t. } \int_{\mathbb{R}^3} \frac{|\widehat{f}(k)|^2}{|k|^2} dk < +\infty \right\}.$$

Then, equation (5) has at least one solution P such that

$$P - P_-^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda), P_\pm^0(P - P_-^0)P_\pm^0 \in \mathfrak{S}_1(\mathfrak{H}_\Lambda), \text{ and } \rho_{P-P_-^0} \in \mathcal{C} \cap L^2(\mathbb{R}^3). \quad (6)$$

All such solutions share the same density $\rho_{P-P_-^0}$.

In (6), $\mathfrak{S}_1(\mathfrak{H}_\Lambda)$ and $\mathfrak{S}_2(\mathfrak{H}_\Lambda)$ are respectively the spaces of trace-class and Hilbert-Schmidt operators [30] on \mathfrak{H}_Λ , and $P_+^0 = 1 - P_-^0$. The method used in [16, 13] was to identify solutions of (5) with minimizers of the so-called reduced Bogoliubov-Dirac-Fock energy which is nothing but the formal difference between the reduced Hartree-Fock energy of P and that of the reference state P_-^0 . Note that due to the uniqueness of $\rho_{P-P_-^0}$ the mean-field operator D is also unique and only δ can differ between two solutions of (5).

Let us mention that it is natural to look for a solution of (5) such that $P - P_-^0$ is a Hilbert-Schmidt operator on \mathfrak{H}_Λ . If P is a projector, the Shale-Stinespring theorem [29] then tells us that P yields a Fock representation equivalent to that of P_-^0 . Even when P is not a projector, it will be associated with a unique Bogoliubov mixed state in the Fock space representation of P_-^0 . This is a mathematical formulation of the statement that P should not be too far from P_-^0 . Indeed, if P is an orthogonal projector, one has (see [15, Lemma 2] and [17, Lemma 1])

$$\left. \begin{array}{l} P - P_-^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda) \\ P^2 = P \end{array} \right\} \implies P_\pm^0(P - P_-^0)P_\pm^0 \in \mathfrak{S}_1(\mathfrak{H}_\Lambda), \text{ and } \rho_{P-P_-^0} \in \mathcal{C} \cap L^2(\mathbb{R}^3),$$

therefore, in this case, (6) is just equivalent to the Shale-Stinespring condition $P - P_-^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$.

The property $P_\pm^0(P - P_-^0)P_\pm^0 \in \mathfrak{S}_1(\mathfrak{H}_\Lambda)$ allows us to define the total charge of the system by (see [15])

$$\text{tr}_{P_-^0}(P - P_-^0) := \text{tr}(P_-^0(P - P_-^0)P_-^0) + \text{tr}(P_+^0(P - P_-^0)P_+^0).$$

When P is a projector, the above quantity is always an integer which is indeed nothing but the relative index of the pair (P, P_-^0) (see [15, 1]). Varying μ allows to pick the desired total charge. In particular, if ν is small enough and $\mu = 0$, then one has $\|P - P_-^0\| < 1$ and the relative index vanishes

$$\text{tr}_{P_-^0}(P - P_-^0) = 0.$$

It is very important to realize that solutions of (5) are singular mathematical objects. This fact is precisely at the origin of charge renormalization. In [13, Theorem 1], the following was proved.

Theorem 2 (Nonperturbative charge renormalization formula [13]). *Assume that $\alpha \geq 0$, $\Lambda > 0$ and $\mu \in (-1, 1)$ are given. If $\nu \in \mathcal{C} \cap L^1(\mathbb{R}^3)$, then $\rho_{P-P_-^0} \in L^1(\mathbb{R}^3)$ and*

$$\int_{\mathbb{R}^3} \nu - \int_{\mathbb{R}^3} \rho_{P-P_-^0} = \frac{\int_{\mathbb{R}^3} \nu - \text{tr}_{P_-^0}(P - P_-^0)}{1 + \alpha B_\Lambda}. \quad (7)$$

Remark 1. A related model was studied by Cancès and Lewin [3] for defects in non-relativistic crystals. In contrast to the Dirac case, the charge density of the polarized Fermi sea is not necessarily integrable. In particular it is not in $L^1(\mathbb{R}^3)$ for anisotropic materials and the question is open in the isotropic case.

In Formula (7), B_Λ is an explicit function of the ultraviolet cut-off Λ (see the comments after (2.4) and (B.10)), which behaves like

$$B_\Lambda = \frac{2}{3\pi} \log \Lambda - \frac{5}{9\pi} + \frac{2 \log 2}{3\pi} + O\left(\frac{1}{\Lambda^2}\right).$$

Let us emphasize that (7) is non perturbative and holds for all $\alpha \geq 0$ and all $\mu \in (-1, 1)$. Theorem 2 shows that the operator $P - P_-^0$ is in general not trace-class. If $P - P_-^0 \in \mathfrak{S}_1(\mathfrak{H}_\Lambda)$, then it must hold $\text{tr}_{P_-^0}(P - P_-^0) = \text{tr}(P - P_-^0) = \int_{\mathbb{R}^3} \rho_{P-P_-^0}$.

In our model we have two possible definitions of the charge of the system: $\int_{\mathbb{R}^3} \nu - \text{tr}_{P_-^0}(P - P_-^0)$ and $\int_{\mathbb{R}^3} (\nu - \rho_{P-P_-^0})$. In practice it is the electrostatic field induced by the nucleus (together with the vacuum polarization density) which is measured, hence it is more natural to define the charge by means of the density. By (7), the total Coulomb potential is, at infinity,

$$\alpha(\nu - \rho_{P-P_-^0}) * \frac{1}{|x|} \underset{|x| \rightarrow +\infty}{\sim} \alpha \frac{\int_{\mathbb{R}^3} (\nu - \rho_{P-P_-^0})}{|x|} = \frac{\alpha}{1 + \alpha B_\Lambda} \left(\frac{\int_{\mathbb{R}^3} \nu - \text{tr}_{P_-^0}(P - P_-^0)}{|x|} \right).$$

Let us assume for simplicity that we put in the vacuum ($\mu = 0$) a nucleus containing $\int_{\mathbb{R}^3} \nu = Z$ protons and which is small enough in the sense that $\|\nu\|_C \ll 1$. Then $\text{tr}_{P_-^0}(P - P_-^0) = 0$ by [16, Theorem 3] and we see that at infinity the potential induced by the nucleus is not $\alpha Z/|x|$ as expected, but rather $\alpha_{\text{ph}} Z/|x|$ where

$$\alpha_{\text{ph}} = Z_3 \alpha \text{ with } Z_3 = \frac{1}{1 + \alpha B_\Lambda} = 1 - \alpha_{\text{ph}} B_\Lambda. \quad (8)$$

Our charge renormalization constant ² Z_3 agrees with the one appearing in the one-loop renormalization of full QED [9, 2, 21]. In our model it allows to renormalize the density of charge to all orders, as we will explain in Theorem 3. The value of α is not observable, α_{ph} is the real physical constant since we always observe the nucleus together with the vacuum polarization density. Its experimental value is $\alpha_{\text{ph}} \simeq 1/137$.

In our theory we must therefore fix α_{ph} and not α . Using (8) we can express any physical quantity in terms of α_{ph} and Λ only. Unfortunately it holds $\alpha_{\text{ph}} B_\Lambda < 1$ hence it makes no sense to take $\Lambda \rightarrow +\infty$ while keeping α_{ph} fixed (this is the so-called Landau pole [22, 23]) and one has to look for a weaker definition of renormalizability. The cut-off Λ which was first introduced as a mathematical trick to regularize the model has actually a physical meaning. Because of the above constraint $\alpha_{\text{ph}} B_\Lambda < 1$, a natural scale occurs beyond which the model does not make sense. Fortunately, this scale is of the order $e^{3\pi/2\alpha_{\text{ph}}}$, a huge number for $\alpha_{\text{ph}} \simeq 1/137$.

It is more convenient to change variables and take as new parameters α_{ph} and $Z_3 = 1 - \alpha_{\text{ph}} B_\Lambda$, with the additional constraint that $0 < Z_3 < 1$. The new parameter Z_3 is now independent of α_{ph} and the natural question arises whether predicted physical quantities will depend very much on the chosen value of $0 < Z_3 < 1$. The purpose of this paper is to prove that the asymptotics of any physical quantity in the regime $\alpha_{\text{ph}} \ll 1$ is actually independent of Z_3 to any order in α_{ph} , which is what we call asymptotic renormalizability. Note that fixing $Z_3 \in (0, 1)$ amounts to take $\Lambda \simeq C e^{3\pi(1-Z_3)/2\alpha_{\text{ph}}} \gg 1$.

Instead of looking at all possible physical observables, it is convenient to define a renormalized density ρ_{ph} . Following [16], we define it by the relation

$$\alpha_{\text{ph}} \rho_{\text{ph}} = \alpha(\nu - \rho_{P-P_-^0}), \quad (9)$$

²The renormalization constant Z_3 should not be confused with the nuclear charge $Z = \int_{\mathbb{R}^3} \nu$.

in such a way that $D = D^0 - \alpha_{\text{ph}}\rho_{\text{ph}} * |x|^{-1}$. This procedure is similar to wave-function renormalization. By uniqueness of $\rho_{P-P^0_-}$ we can see ρ_{ph} as a function of α_{ph} , ν , μ and Λ (or Z_3). For the sake of clarity we will not emphasize the dependence in ν and μ which will be fixed quantities. Also we will use the same notation $\rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda)$ or $\rho_{\text{ph}}(\alpha_{\text{ph}}, Z_3)$, depending on the context. The self-consistent equation for ρ_{ph} was derived in [16] and it is mentioned below in Section 2.

From now on, we will assume that

$$\mu = 0.$$

For small external densities ν , this means that we will be looking at the vacuum polarization in the presence of the nucleus, without considering any real electron (that is, ρ_{ph} is the renormalized density of the nucleus containing both the bare density ν and the vacuum polarization density $\rho_{P-P^0_-}$). We will explain in Section 2 that one can expand $\rho_{\text{ph}} = \rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda)$ as follows

$$\rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda) = \sum_{n=0}^{+\infty} (\alpha_{\text{ph}})^n \nu_{n,\Lambda}, \quad (10)$$

where $(\nu_{n,\Lambda})_{n \in \mathbb{N}} \in (L^2(\mathbb{R}^3) \cap \mathcal{C})^{\mathbb{N}}$ is a sequence depending only on the external density ν and the cut-off Λ . This sequence is defined below in Section 2. The series (10) has a positive radius of convergence, which is however believed to shrink to zero when $\Lambda \rightarrow +\infty$.

Assuming $\widehat{\nu}$ decays fast enough (see condition (11)), we will prove that for any fixed n , the limit $\nu_{n,\Lambda} \rightarrow \nu_n$ exists in $L^2(\mathbb{R}^3) \cap \mathcal{C}$. This is what is usually meant by renormalizability in QED. Each term of the perturbation series in powers of the physical α_{ph} has a limit when the cut-off is removed. The sequence $(\nu_n)_{n \in \mathbb{N}}$ is the one which is calculated in practice [2, 14, 12, 11]. One has for instance $\nu_0 = \nu$ and

$$\nu_1 * \frac{1}{|x|} = \frac{1}{3\pi} \int_1^{+\infty} (t^2 - 1)^{\frac{1}{2}} \left(\frac{2}{t^2} + \frac{1}{t^4} \right) \left(\int_{\mathbb{R}^3} e^{-2|x-y|t} \frac{\nu(y)}{|x-y|} dy \right) dt,$$

the Uehling potential [33, 28]. All the other ν_n can be calculated by induction in terms of ν_0, \dots, ν_{n-1} , as is explained below in Section 2.

The next natural question is to understand the link between the well-defined, cut-off dependent, series (10) and the formal series $\sum_{n=0}^{+\infty} (\alpha_{\text{ph}})^n \nu_n$. Recall that $\alpha_{\text{ph}} B_\Lambda < 1$ by construction, so it is in principle not allowed to take the limit $\Lambda \rightarrow +\infty$ while keeping α_{ph} fixed. We rather want to think of $Z_3 = 1 - \alpha_{\text{ph}} B_\Lambda$ as being fixed. The main result in this paper is the following

Theorem 3 (Asymptotic renormalization of the nuclear charge density). *Consider a function $\nu \in L^2(\mathbb{R}^3) \cap \mathcal{C}$ such that*

$$\int_{\mathbb{R}^3} \log(1 + |k|)^{2N+2} |\widehat{\nu}(k)|^2 dk < +\infty, \quad (11)$$

for some integer N . Let $\rho_{\text{ph}}(\alpha_{\text{ph}}, Z_3)$ be the unique physical density defined by (9) with $\mu = 0$, $Z_3 = 1 - \alpha_{\text{ph}} B_\Lambda$ and $\alpha_{\text{ph}} = Z_3 \alpha$.

Then, for every $0 < \epsilon < 1/2$, there exist two constants $C(N, \epsilon, \nu)$ and $a(N, \epsilon, \nu)$, depending only on N , ϵ and ν , such that one has

$$\left\| \rho_{\text{ph}}(\alpha_{\text{ph}}, Z_3) - \sum_{n=0}^N \nu_n (\alpha_{\text{ph}})^n \right\|_{L^2(\mathbb{R}^3) \cap \mathcal{C}} \leq C(N, \epsilon, \nu) \alpha_{\text{ph}}^{N+1}, \quad (12)$$

for all $0 \leq \alpha_{\text{ph}} \leq a(N, \epsilon, \nu)$ and all $\epsilon \leq Z_3 \leq 1 - \epsilon$.

Remark 2. We will provide explicit induction formulas for the sequence $(\nu_n)_{n \in \mathbb{N}}$ later in Section 2. In particular, we will see in the proof that under Assumption (11), one has $\nu_n \in L^2(\mathbb{R}^3) \cap \mathcal{C}$ for all $0 \leq n \leq N$. Therefore the approximation series of order N appearing in (12), $\sum_{n=0}^N (\alpha_{\text{ph}})^n \nu_n$, is a well-defined function of $L^2(\mathbb{R}^3) \cap \mathcal{C}$.

Remark 3. The space $L^2(\mathbb{R}^3) \cap \mathcal{C}$ is the natural space which occurs in this theory. In particular the Coulomb norm is nothing but the classical electrostatic energy which appears in the reduced BDF energy functional. Our result can be extended to Sobolev spaces $H^s(\mathbb{R}^3)$ provided ν is smooth enough.

The interpretation of Theorem 3 is that the renormalized density $\rho_{\text{ph}}(\alpha_{\text{ph}}, Z_3)$ is asymptotically (meaning up to any fixed order N) given by the formal series $\sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_n$, uniformly in the renormalization parameter Z_3 in the range $\epsilon \leq Z_3 \leq 1 - \epsilon$. Therefore, for a very large range of cut-offs, essentially

$$C_1 e^{\frac{3\epsilon\pi}{2\alpha_{\text{ph}}}} \leq \Lambda \leq C_2 e^{\frac{3(1-\epsilon)\pi}{2\alpha_{\text{ph}}}},$$

the result is independent of Λ and it is given by the formal series $\sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_n$. Our formulation of renormalizability is more precise than the requirement that each $\nu_{n,\Lambda}$ converges. It also leads to the formal perturbation series in a very natural way.

A natural question is to ask for the convergence of the perturbation series $\sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_n$. It was argued by Dyson in [10] that it is probably divergent, but we are unable to transform his argument into a rigorous mathematical proof. We will make more comments on the series $\sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_n$ at the end of next section.

It would be interesting to extend Theorem 3 to the case of atoms with 'real' electrons. This amounts to taking μ sufficiently close to 1 at the same time as α_{ph} is small. However this case is more difficult than what is done here: an additional expansion of the electronic charge density in powers of α_{ph} is needed.

Another interesting problem is to include the exchange term in the model. As for the existence of solutions, this was already done for the free vacuum in [24, 19] and in an external field in [15, 19, 17]. In this case the mass of the electron also has to be renormalized [19, Sec. 2.5] and there is probably no explicit formula for the corresponding m_{ph} . In fact, it is not clear at all whether the renormalization program can be applied: In [26, pp. 194-195], it is argued that mass and charge renormalization alone are not enough to completely remove the divergences by means of multiplicative parameters, in mean-field theory with exchange term.

The proof of Theorem 3 (given in Section 5 below) is divided into two steps. We first estimate the difference (see Lemma 5.1)

$$\left\| \rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda) - \sum_{n=0}^N \nu_{n,\Lambda} (\alpha_{\text{ph}})^n \right\|_{L^2(\mathbb{R}^3) \cap \mathcal{C}} \leq C_1(N, \epsilon, \nu) (\alpha_{\text{ph}})^{N+1} \quad (13)$$

for a constant $C_1(N, \epsilon, \nu)$ depending only on N , ϵ and ν , and under the assumption that $\epsilon \leq Z_3 = 1 - \alpha_{\text{ph}} B_\Lambda \leq 1 - \epsilon$. This amounts to expanding the solution of the self-consistent equation (5) up to the N th order in α_{ph} while controlling the error term uniformly in Λ . Then we show in Lemma 5.2 that

$$\forall 0 \leq n \leq N, \|\nu_{n,\Lambda} - \nu_n\|_{L^2(\mathbb{R}^3) \cap \mathcal{C}} \leq \frac{C_2(N, \nu)}{(B_\Lambda)^{N+1-n}}, \quad (14)$$

for a constant $C_2(N, \nu)$ depending only on N and ν , leading to the bound

$$\left\| \sum_{n=0}^N \nu_{n,\Lambda} (\alpha_{\text{ph}})^n - \sum_{n=0}^N \nu_n (\alpha_{\text{ph}})^n \right\|_{L^2(\mathbb{R}^3) \cap \mathcal{C}} \leq (\alpha_{\text{ph}})^{N+1} \frac{C_2(N, \nu)(1 - \epsilon^{N+1})}{\epsilon^{N+1}(1 - \epsilon)}, \quad (15)$$

since by assumption $(B_\Lambda)^{-N-1+n} \leq (\alpha_{\text{ph}}/\epsilon)^{N+1-n}$. The main result then follows from (13) and (15). All these bounds strongly use the explicit recursion relations defining the sequences $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ and $(\nu_n)_{n \in \mathbb{N}}$, as well as tedious estimates on the nonlinear terms appearing in these relations.

The rest of the paper is organized as follows. In Section 2 we define the sequences $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ and $(\nu_n)_{n \in \mathbb{N}}$ by their respective recursion formulas and we discuss some properties of the latter. In particular, in Theorem 2.1, we give a simple estimate on $\|\nu_n\|_{L^2(\mathbb{R}^3) \cap \mathcal{C}}$. In Section 3 we present estimates on the different terms appearing in the recursion formulas. Of particular interest will be the density $\nu_{1,\Lambda}$ giving rise to the Uehling potential. In Fourier space, we have $\widehat{\nu}_{1,\Lambda}(k) = U_\Lambda(k)\widehat{\nu}(k)$ for an explicit function $U_\Lambda(k)$ which is studied in Section 3.1. The proofs of Theorems 2.1 and 3 are respectively provided in Sections 4 and 5. Some other technical proofs are provided in Appendices A, B and C.

Acknowledgements. The authors are grateful to Christian Brouder for interesting comments. M.L. would like to thank Jan Dereziński and Jan Philip Solovej for stimulating discussions. Grants from the French Ministry of Research (M.L. and É.S., ANR-10-BLAN 0101) and from the European Research Council (M.L., MNIQS-258023) are gratefully acknowledged.

2 The two sequences $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ and $(\nu_n)_{n \in \mathbb{N}}$

In this section we derive formulas for $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ and $(\nu_n)_{n \in \mathbb{N}}$, and we make some comments on the latter.

2.1 Definition of $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ and $(\nu_n)_{n \in \mathbb{N}}$

We start with the self-consistent equation (5) with cut-off, assuming $\mu = 0$. Note that in the regime of interest in Theorem 3, we have $\alpha = \alpha_{\text{ph}}/Z_3 \leq \alpha_{\text{ph}}/\epsilon$. When $\alpha\pi^{1/6}2^{11/6}\|\nu\|_{\mathcal{C}} < 1$, it is known that $0 \notin \sigma(D)$ hence $\delta = 0$ in (5) (see [16, Theorem 3] and [13, Lemma 11]). Therefore assuming $a(\nu, N, \epsilon) \leq \epsilon(\pi^{1/6}2^{11/6}\|\nu\|_{\mathcal{C}})^{-1}$ in Theorem 3, we automatically have that $\delta = 0$ and $P = P^2$ is unique.

The idea is then to expand the self-consistent equation

$$P = \chi_{(-\infty, 0)} \Pi_\Lambda \left(D^0 + \alpha(\rho_{P-P^0} - \nu) * \frac{1}{|x|} \right) \Pi_\Lambda, \quad (2.1)$$

in powers of α by means of the resolvent formula. A similar computation was performed in [20] in the Furry picture (that is, neglecting the self-consistent terms). This method was also used in [15] to prove existence and uniqueness of solutions of the self-consistent equation. We define

$$F_{n,\Lambda}(\mu_1, \dots, \mu_k) := \rho \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{D^0 + i\eta} \prod_{j=1}^n \left(\Pi_\Lambda \left(\mu_j * \frac{1}{|x|} \right) \Pi_\Lambda \frac{1}{D^0 + i\eta} \right) d\eta \right), \quad (2.2)$$

where we recall that Π_Λ is the orthogonal projector onto \mathfrak{H}_Λ in $L^2(\mathbb{R}^3, \mathbb{C}^4)$ and $(\mu_1, \dots, \mu_n) \in \mathcal{C}^n$. We will always use the simplified notation $F_{n,\Lambda}(\mu) := F_{n,\Lambda}(\mu, \dots, \mu)$ and $\nu_\Lambda := \mathcal{F}^{-1}(\widehat{\nu} \mathbb{1}_{B(0, 2\Lambda)})$. Note that by Furry's theorem $F_{2j,\Lambda} \equiv 0$ for all j (see [15, p. 547]). We also introduce

$$F_\Lambda(\mu) := \sum_{n \geq 3} F_{n,\Lambda}(\mu).$$

The self-consistent equation (2.1) may then be written in terms of the density in Fourier space [15, 16], as

$$\widehat{\rho_{P-P^0}}(k) = -\alpha B_\Lambda(k) (\widehat{\rho_{P-P^0}}(k) - \widehat{\nu}_\Lambda(k)) + \widehat{F}_\Lambda(\alpha(\nu - \rho_{P-P^0})), \quad (2.3)$$

where the function $B_\Lambda(k)$ is given by

$$B_\Lambda(k) = \frac{1}{\pi} \int_0^{Z_\Lambda(|k|)} \frac{z^2 - \frac{z^4}{3}}{(1-z^2)(1 + \frac{|k|^2(1-z^2)}{4})} dz + \frac{|k|}{2\pi} \int_0^{Z_\Lambda(|k|)} \frac{z - \frac{z^3}{3}}{\sqrt{1+\Lambda^2 - \frac{|k|z}{2}}} dz, \quad (2.4)$$

with $Z_\Lambda(r) = (\sqrt{1+\Lambda^2} - \sqrt{1+(\Lambda-r)^2})/r$ (see [13]). The formula for $B_\Lambda(k)$ is well-known (but in most previous works the second term was ignored (see for instance [25])).

Defining $U_\Lambda(|k|) = B_\Lambda - B_\Lambda(k)$ where $B_\Lambda = B_\Lambda(0)$ and $0 \leq U_\Lambda(|k|) \leq B_\Lambda$ with $U_\Lambda(2\Lambda) = B_\Lambda$, we get the renormalized equation

$$(1 - \alpha_{\text{ph}} U_\Lambda) \widehat{\rho}_{\text{ph}} + \widehat{F}_\Lambda(\alpha_{\text{ph}} \rho_{\text{ph}}) = \widehat{\nu}_\Lambda, \quad (2.5)$$

with the renormalized coupling constant $\alpha_{\text{ph}} := \alpha/(1 + \alpha B_\Lambda)$ and the renormalized density $\alpha_{\text{ph}} \rho_{\text{ph}} = \alpha(\nu - \rho_Q)$ (see [16]). For convenience, we will denote by \mathcal{U}_Λ the operator of multiplication by the function $U_\Lambda(|k|)$ in the Fourier domain. Hence we can write the self-consistent equation (2.5) in direct space as

$$(1 - \alpha_{\text{ph}} \mathcal{U}_\Lambda) \rho_{\text{ph}} + F_\Lambda(\alpha_{\text{ph}} \rho_{\text{ph}}) = \nu_\Lambda. \quad (2.6)$$

We now expand the unique solution $\rho_{\text{ph}} = \rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda)$ of (2.6) in powers of α_{ph} . Writing a formal series

$$\rho_{\text{ph}} = \sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_{n,\Lambda}, \quad (2.7)$$

we find that the functions $\nu_{n,\Lambda}$ must satisfy the following recurrence relation

$$\begin{cases} \nu_{0,\Lambda} = \nu_\Lambda, \\ \nu_{1,\Lambda} = \mathcal{U}_\Lambda \nu_\Lambda, \\ \nu_{n,\Lambda} = \mathcal{U}_\Lambda \nu_{n-1,\Lambda} + \sum_{j=3}^n \sum_{n_1+\dots+n_j=n-j} F_{j,\Lambda}(\nu_{n_1,\Lambda}, \dots, \nu_{n_j,\Lambda}), \quad \forall n \geq 2. \end{cases} \quad (2.8)$$

Note that the operator \mathcal{U}_Λ is bounded by $U(2\Lambda) = B_\Lambda$ on \mathfrak{H}_Λ and that, as we will see later in Corollary 3.1, each $F_{j,\Lambda}$ is continuous on \mathcal{C}^j with values in $L^2 \cap \mathcal{C}$. The sequence $(\nu_{n,\Lambda})_{n \in \mathbb{N}}$ is thus well-defined in $L^2 \cap \mathcal{C}$. Using estimates from [15] it can be proven that the series (2.7) has a finite radius of convergence in $L^2 \cap \mathcal{C}$, but this is not needed for the moment and we can stay at a formal level in this section.

We can now formally pass to the limit as $\Lambda \rightarrow +\infty$ and define by induction a sequence $(\nu_n)_{n \in \mathbb{N}}$ by

$$\begin{cases} \nu_0 = \nu, \\ \nu_1 = \mathcal{U} \nu, \\ \nu_n = \mathcal{U} \nu_{n-1} + \sum_{j=3}^n \sum_{n_1+\dots+n_j=n-j} F_j(\nu_{n_1}, \dots, \nu_{n_j}), \quad \forall n \geq 2. \end{cases} \quad (2.9)$$

where the F_j are defined similarly as the $F_{j,\Lambda}$ with Π_Λ removed and \mathcal{U} is the operator of multiplication by the function $U(|k|)$ in the Fourier domain, defined by

$$\begin{aligned} U(r) &:= \lim_{\Lambda \rightarrow +\infty} U_\Lambda(r) = \frac{r^2}{4\pi} \int_0^1 \frac{z^2 - \frac{z^4}{3}}{1 + \frac{r^2(1-z^2)}{4}} dz \\ &= \frac{12 - 5r^2}{9\pi r^2} + \frac{\sqrt{4+r^2}}{3\pi r^3} (r^2 - 2) \log \left(\frac{\sqrt{4+r^2} + r}{\sqrt{4+r^2} - r} \right). \end{aligned} \quad (2.10)$$

If the series $\sum_{n \geq 0} \nu_n(\alpha_{\text{ph}})^n$ had a positive radius of convergence, it would be a solution of the renormalized equation without ultraviolet cut-off

$$(1 - \alpha_{\text{ph}}\mathcal{U})\rho_{\text{ph}} + F(\alpha_{\text{ph}}\rho_{\text{ph}}) = \nu. \quad (2.11)$$

However, it is widely believed that the series is divergent (as discussed in the next subsection) and it is not known whether the renormalized equation (2.11) admits any solution, even for α_{ph} small enough.

2.2 On the series $\sum_{n \geq 0} \nu_n(\alpha_{\text{ph}})^n$

The recursion formula (2.9) defining $(\nu_n)_{n \in \mathbb{N}}$ contains two terms. The first term $\mathcal{U}\nu_{n-1}$ is a simple multiplication operator in Fourier space, by the function $U(|k|)$ which diverges at infinity. The second term involves the nonlinear functions F_j 's. If only the first term with \mathcal{U} were present, the series $\sum_{n \geq 0} \nu_n(\alpha_{\text{ph}})^n$ would only converge when the Fourier transform $\widehat{\nu}$ has a compact support, the radius of convergence depending on the size of this support. If only the nonlinear terms were present, the series would have a finite radius of convergence by the estimates of [15] and of Section 3.2.

However when the two terms are combined, the situation is much more complicated. The nonlinear terms act like convolutions in Fourier space, hence even if $\widehat{\nu}$ has a compact support in the Fourier domain, the support of $\widehat{\nu}_n$ will probably grow with n . A careful study of the mixed effect of the multiplication by the divergent function U and the nonlinearities seems rather difficult. We will prove the following estimate:

Theorem 2.1 (Estimate on $(\nu_n)_{n \geq 1}$). *There exist universal constants A and K such that*

$$\|(1 + \mathcal{U})^{m-n}\nu_n\|_{L^2 \cap \mathcal{C}} \leq A^{n+1} \max \left\{ \|(1 + \mathcal{U})^m\nu\|_{L^2 \cap \mathcal{C}}, (K \log(m))^{\frac{mn}{2}} \|(1 + \mathcal{U})^m\nu\|_{L^2 \cap \mathcal{C}}^{n+1} \right\}, \quad (2.12)$$

for all $\Lambda \geq 1$, $m \in \mathbb{N}$ and $0 \leq n \leq m$.

Even if we assume that ν decays fast enough in Fourier space, for instance

$$\forall n \geq 0, \|(1 + \mathcal{U})^n\nu\|_{L^2 \cap \mathcal{C}} \leq C^n,$$

the above estimate (2.12) does not imply that the series $\sum_{n \geq 0} \nu_n(\alpha_{\text{ph}})^n$ is convergent for α_{ph} small enough. Although our estimate (2.12) is certainly far from optimal, as we have already mentioned, it is expected that the series does not converge in any appropriate sense [10].

It is sometimes argued that the series could be Borel summable. The Borel transform is defined by

$$\mathcal{B}(t) = \sum_{n \geq 0} \frac{t^n}{n!} \nu_n.$$

If $\mathcal{B}(t)$ is a convergent series (for an appropriate norm) having a holomorphic extension to a domain containing the positive real line, such that

$$\tilde{\mathcal{B}}(\alpha_{\text{ph}}) := \int_0^{+\infty} \mathcal{B}(t) e^{-\frac{t}{\alpha_{\text{ph}}}} dt$$

makes sense in an appropriate neighborhood of $\alpha_{\text{ph}} = 0$, one may see $\tilde{\mathcal{B}}(\alpha_{\text{ph}})$ as the physical density, whose series $\sum_{n \geq 0} (\alpha_{\text{ph}})^n \nu_n$ is only asymptotic. Proving such results mathematically is hard, even for the model studied in this paper. Our estimate (2.12) does not even allow to define the Borel transform $\mathcal{B}(t)$ in $L^2(\mathbb{R}^3) \cap \mathcal{C}$.

But Borel summability is not the only tool to construct a physical density providing the correct asymptotic series. For the model studied in the present paper, we have several natural families of functions of α_{ph} : the cut-off densities

$$\rho_{\text{ph}}\left(\alpha_{\text{ph}}, Ce^{\frac{3(1-Z_3)\pi}{2\alpha_{\text{ph}}}}\right) \quad (2.13)$$

obtained by minimizing the reduced BDF energy with a cut-off $\Lambda = Ce^{3(1-Z_3)\pi/2\alpha_{\text{ph}}}$ and using the relation (9). Each such density (2.13) has (for fixed C and $0 < Z_3 < 1$) the required asymptotic series in α_{ph} by Theorem 3, and it solves the self-consistent equation (5) with the corresponding cut-off Λ . Furthermore this solution has the benefit of being well-defined even when α_{ph} is not small, allowing for the description of nonperturbative physical events.

The rest of the paper is devoted to the proofs of Theorems 3 and 2.1.

3 Some preliminary results

In this section we state two preliminary results that will be useful in the proof of our main results, Theorems 3 and 2.1. The corresponding lengthy calculations will be provided later in Appendices A, B and C.

Notation. In the whole paper we use the notation $E(r) = (1 + \|r\|^2)^{1/2}$ for $r \in \mathbb{R}^3$ or $r \in \mathbb{R}$.

3.1 The Uehling multiplier U

The operator \mathcal{U} , defined previously as the multiplication by the function U in the Fourier domain, plays a major role in the definition of the sequence $(\nu_n)_{n \in \mathbb{N}}$. In this section, we provide precise estimates quantifying the convergence of U_Λ towards U when $\Lambda \rightarrow +\infty$, which will be very useful in the proof of Theorem 3.

Proposition 3.1. *Let $\Lambda \geq 1$ and denote by U_Λ , the function defined on \mathbb{R}_+ by*

$$U_\Lambda(r) = \begin{cases} B_\Lambda - B_\Lambda(r), & \text{when } 0 \leq r \leq 2\Lambda, \\ 0, & \text{otherwise.} \end{cases} \quad (3.1)$$

Then, for all $r \in \mathbb{R}_+$ it holds $\lim_{\Lambda \rightarrow +\infty} U_\Lambda(r) = U(r)$. Moreover, for $\kappa_0 = 15\pi/2$,

$$\forall m \geq 0, \left\| \frac{U_\Lambda - U}{(1 + U)^{m+1}} \right\|_{L^\infty} \leq \kappa_0^{m+3} \max \left\{ \frac{1}{(1 + B_\Lambda)^m}, \frac{1}{E(2\Lambda)} \right\}. \quad (3.2)$$

Finally, one has for a universal constant κ_1 (given in Lemma B.1 below)

$$\forall 0 \leq r \leq 2\Lambda, 0 \leq U_\Lambda(r) \leq \kappa_1(1 + U(r)). \quad (3.3)$$

Proposition 3.1 is proved in Appendix B. Note that the uniform estimate (3.2) will later yield our estimate (14) on $\nu_{n,\Lambda} - \nu_n$ (see Lemma 5.2). More properties of U and U_Λ are provided later in Appendix A.

3.2 The nonlinear terms $F_{n,\Lambda}$ and F_n

In this section, we provide estimates on the functions $F_{n,\Lambda}$ and F_n , which will be one of the main ingredients in the proof of Theorem 3. We recall that $F_{2n,\Lambda} = F_{2n} = 0$ by Furry's theorem

(see [15, p. 547]). In order to state our main result, we introduce the functions

$$F_{n,\Lambda}^\epsilon(\mu) = \rho \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{D^0 + i\eta} \prod_{j=1}^n \left(\Pi_\Lambda^{(\epsilon_j)} \mu_j * \frac{1}{|x|} \Pi_\Lambda^{(\epsilon_{j+1})} \frac{1}{D^0 + i\eta} \right) d\eta \right), \quad (3.4)$$

for any $n \geq 3$, $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{C}^n$ and $\epsilon = (\epsilon_1, \dots, \epsilon_{n+1}) \in \{-1, 0, 1\}^{n+1}$. Here, we have used the notation

$$\Pi_\Lambda^{(1)} := \Pi_\Lambda, \quad \Pi_\Lambda^{(-1)} := 1 - \Pi_\Lambda \quad \text{and} \quad \Pi_\Lambda^{(0)} := 1 = \Pi_\Lambda^{(1)} + \Pi_\Lambda^{(-1)}. \quad (3.5)$$

The main result of this section is the following

Proposition 3.2 (Estimates on $F_{n,\Lambda}^\epsilon$). *Let $m \in \mathbb{N}$, $\Lambda \geq 1$ and $\epsilon \in \{-1, 0, 1\}^{n+1}$. Assume that $n \geq 3$. Then, there exist universal constants C and K such that*

$$\|(1 + \mathcal{U})^m F_{n,\Lambda}^\epsilon(\mu)\|_{L^2 \cap \mathcal{C}} \leq \frac{C^n (K \log n)^m}{\Lambda^{\frac{n(\epsilon)}{24}}} \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (3.6)$$

for all $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{C}^n$. Here, $n(\epsilon) = 1$, if at least one ϵ_j is equal to -1 , and $n(\epsilon) = 0$ otherwise.

By (2.2), (3.4) and (3.5), we can write $F_{n,\Lambda}^{(1,\dots,1)} = F_{n,\Lambda}$ and $F_{n,\Lambda}^{(0,\dots,0)} = F_n$. Therefore the following is a byproduct of (3.6):

Corollary 3.1. *Let $m \geq 0$, $\Lambda \geq 1$ and $n \geq 3$, an odd integer. Then,*

$$\max \left\{ \|(1 + \mathcal{U})^m F_{n,\Lambda}(\mu)\|_{L^2 \cap \mathcal{C}}, \|(1 + \mathcal{U})^m F_n(\mu)\|_{L^2 \cap \mathcal{C}} \right\} \leq C^m (K \log n)^m \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (3.7)$$

for any $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{C}^n$. Here, C and K refer to the universal constants given by Proposition 3.2. In particular, the functions $F_{n,\Lambda}$ and F_n are continuous on \mathcal{C}^n with values in $L^2 \cap \mathcal{C}$.

Recall $F_{2k} = F_{2k,\Lambda} = 0$ hence only the case of n being an odd integer is relevant. The estimates of Proposition 3.2 are an adaptation of ideas of [15], in which similar bounds were computed (see e.g. Lemmas 15 and 16 in [15]). Notice however that the projector Π_Λ was never mentioned in [15] since Λ was a fixed number. We focus here on the limit $\Lambda \rightarrow +\infty$ and we need to quantify the dependence on Λ of the estimates on the functions $F_{n,\Lambda}^\epsilon$. The proof of Proposition 3.2 is provided below in Appendix C. The factor $(K \log n)^m$ comes from (A.7) of Lemma A.3 and the constant K is also the one appearing in Theorem 2.1.

4 Proof of Theorem 2.1

This section is devoted to the proof of our estimate (2.12) on the n th order density ν_n . The definition of $\nu_{n,\Lambda}$ being very similar to that of ν_n , our proof also provides the following

Proposition 4.1 (Estimates on $\nu_{n,\Lambda}$). *There exists $A > 0$ such that*

$$\|(1 + \mathcal{U})^{m-n} \nu_{n,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq A^{n+1} \max \left\{ \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}, (K \log(m))^{\frac{mn}{2}} \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}^{n+1} \right\}, \quad (4.1)$$

for any $\Lambda \geq 1$, $m \in \mathbb{N}$ and $0 \leq n \leq m$.

We postpone the proof of Proposition 4.1 and first complete that of Theorem 2.1.

Proof of Theorem 2.1. We split the proof into three steps. First, we estimate by means of (3.7), the following norms $J_{m,n} := \|(1 + \mathcal{U})^{m-n}\nu_n\|_{L^2 \cap \mathcal{C}}$.

Step 1. Let $m \in \mathbb{N}$ and denote

$$P_m(t) := \sum_{n=0}^m J_{m,n} t^n.$$

The polynomial $P_m(t)$ satisfies for any $t \geq 0$

$$P_m(t) \leq (1 + t + t^2) \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}} + \mathcal{Q}_m(t P_m(t)), \quad (4.2)$$

where (C and K are the constants of Proposition 3.2)

$$\mathcal{Q}_m(u) := u + \sum_{j=3}^m C^j (K \log j)^{m-j} u^j. \quad (4.3)$$

Let us assume first that $n \in \{0, 1, 2\}$. By (2.9), we then have $\nu_n = \mathcal{U}^n \nu$, hence

$$\forall n \in \{0, 1, 2\}, J_{m,n} = \|(1 + \mathcal{U})^{m-n} \mathcal{U}^n \nu\|_{L^2 \cap \mathcal{C}} \leq \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}. \quad (4.4)$$

We now turn to the case $n \geq 3$. By (2.9), we have

$$\begin{aligned} J_{m,n} &\leq \|(1 + \mathcal{U})^{m-n} \mathcal{U} \nu_{n-1}\|_{L^2 \cap \mathcal{C}} \\ &\quad + \sum_{3 \leq 2j+1 \leq n} \sum_{\sum_{k=1}^{2j+1} n_k = n-2j-1} \|(1 + \mathcal{U})^{m-n} F_{2j+1}(\nu_{n_1}, \dots, \nu_{n_{2j+1}})\|_{L^2 \cap \mathcal{C}}, \end{aligned}$$

hence, by Corollary 3.1,

$$\begin{aligned} J_{m,n} &\leq \|(1 + \mathcal{U})^{m-n+1} \nu_{n-1}\|_{L^2 \cap \mathcal{C}} + \sum_{3 \leq 2j+1 \leq n} \sum_{\sum_{k=1}^{2j+1} n_k = n-2j-1} C^{2j+1} \times \\ &\quad \times (K \log(2j+1))^{m-n} \prod_{k=1}^{2j+1} \|(1 + \mathcal{U})^{m-n} \nu_{n_k}\|_{\mathcal{C}}. \end{aligned}$$

Since $\|(1 + \mathcal{U})^{m-n} \nu_{n_k}\|_{\mathcal{C}} \leq \|(1 + \mathcal{U})^{m-n_k} \nu_{n_k}\|_{L^2 \cap \mathcal{C}}$, we arrive at the inequality

$$J_{m,n} \leq J_{m,n-1} + \sum_{j=3}^n C^j (K \log j)^{m-n} \sum_{\sum_{k=1}^j n_k = n-j} \left(\prod_{k=1}^j J_{m,n_k} \right). \quad (4.5)$$

Combining (4.4) with (4.5), we obtain

$$\begin{aligned} P_m(t) &\leq (1 + t + t^2) \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}} + t P_m(t) \\ &\quad + \sum_{n=3}^m \sum_{j=3}^n C^j t^j (K \log j)^{m-n} \sum_{\sum_{k=1}^j n_k = n-j} \left(\prod_{k=1}^j J_{m,n_k} t^{n_k} \right). \end{aligned}$$

By Fubini's theorem,

$$\begin{aligned} P_m(t) &\leq (1 + t + t^2) \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}} + t P_m(t) \\ &\quad + \sum_{j=3}^m C^j t^j (K \log j)^{m-j} \sum_{p=0}^{m-j} \sum_{\sum_{k=1}^j n_k = p} \left(\prod_{k=1}^j J_{m,n_k} t^{n_k} \right), \quad (4.6) \end{aligned}$$

Noticing that

$$\sum_{p=0}^{m-j} \sum_{\sum_{k=1}^j n_k=p} \left(\prod_{k=1}^j J_{m,n_k} t^{n_k} \right) \leq P_m(t)^j,$$

we deduce (4.2) from (4.6). This completes the proof of Step 1. In the second step of the proof of Theorem 2.1 we compute suitable bounds on \mathcal{Q}_m near the origin.

Step 2. *Let $m \geq 3$. There exists a positive constant $A(C, K)$, depending on C and K , but not on m , such that*

$$\mathcal{Q}_m(u) \leq 2u, \text{ for any } 0 \leq u \leq U_m := \frac{A(C, K)}{(K \log(m))^{\frac{m}{2}}}. \quad (4.7)$$

By the definition (4.3) of \mathcal{Q}_m , we have

$$\mathcal{Q}_m(u) \leq u + \sum_{j=3}^m C^j (K \log(j))^{m-j} u^j \leq u + (K \log(m))^m \sum_{j=3}^m \left(\frac{Cu}{K \log m} \right)^j,$$

hence when $2Cu \leq K \log m$ and $2C^3(K \log(m))^{m-3} u^2 \leq 1$, it holds $\mathcal{Q}_m(u) \leq 2u$. This ends the proof of Step 2.

Step 3. *Conclusion of the proof of Theorem 2.1.*

Since the coefficients $J_{m,n}$ are nonnegative, the function $t \mapsto tP_m(t)$ is either identically equal to 0 (then (2.12) is straightforward), or increasing on \mathbb{R}^+ . In the second case, it tends to $+\infty$ as $t \rightarrow +\infty$, and there exists a unique $T_m > 0$ such that

$$T_m P_m(T_m) = U_m. \quad (4.8)$$

Two situations may then occur. If $T_m \geq 1/4$, by (4.2) and (4.7),

$$P_m(t) \leq 2\|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}} + 2tP_m(t)$$

for all $0 \leq t \leq 1/4$. Hence $P_m(t) \leq 4\|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}$ and

$$J_{m,n} \leq 4^n P_m\left(\frac{1}{4}\right) \leq 4^{n+1} \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}.$$

Otherwise $T_m \leq 1/4$ and in this case we can deduce from (4.2), (4.7) and (4.8) that $U_m/T_m \leq 2\|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}} + 2U_m$. This gives

$$T_m \geq \frac{U_m}{4\|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}}.$$

Combining with (4.8) again, we are led to

$$J_{m,n} \leq \frac{U_m}{T_m^{n+1}} \leq \frac{4^{n+1}}{U_m^n} \|(1 + \mathcal{U})^m \nu\|_{L^2 \cap \mathcal{C}}^{n+1}.$$

Estimate (2.12) then follows from (4.7). □

We now turn to the proof of Proposition 4.1.

Proof of Proposition 4.1. The proof is almost identical. Denoting $J_{m,n}^\Lambda := \|(1+\mathcal{U})^{m-n}\nu_{n,\Lambda}\|_{L^2\cap\mathcal{C}}$ and introducing the polynomial function $P_m^\Lambda(t)$ given by

$$P_m^\Lambda(t) := \sum_{n=0}^m J_{m,n}^\Lambda t^n,$$

we deduce from the definition (2.8), and from (3.3) and (3.7) that

$$P_m^\Lambda(t) \leq (1 + \kappa_1 t + \kappa_1^2 t^2) \|(1+\mathcal{U})^m \nu\|_{L^2\cap\mathcal{C}} + (\kappa_1 - 1)tP_m^\Lambda(t) + \mathcal{Q}_m(tP_m^\Lambda(t)), \quad (4.9)$$

for all $t \geq 0$. Estimate (4.1) then follows by applying to (4.9) the arguments of Steps 2 and 3 of the proof of Theorem 2.1. \square

5 Proof of Theorem 3

This last section is devoted to the proof of our main estimate (12). The proof relies on the identity

$$\rho_{\text{ph}} = \nu_\Lambda + \alpha_{\text{ph}} U_\Lambda \rho_{\text{ph}} - \sum_{3 \leq 2n+1 \leq N} \alpha_{\text{ph}}^{2n+1} F_{2n+1,\Lambda}(\rho_{\text{ph}}, \dots, \rho_{\text{ph}}) - \alpha_{\text{ph}}^{N+1} G_{N+1,\Lambda}, \quad (5.1)$$

where we denote

$$G_{N+1,\Lambda} := \rho \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{D^0 - \alpha_{\text{ph}} \rho_{\text{ph}} * |\cdot|^{-1} + i\eta} \prod_{j=1}^{N+1} \left(\Pi_\Lambda(\rho_{\text{ph}} * \frac{1}{|\cdot|}) \Pi_\Lambda \frac{1}{D^0 + i\eta} \right) d\eta \right). \quad (5.2)$$

The formula (5.1) follows from Cauchy's formula applied to (2.6). As mentioned in the introduction, the proof of (12) naturally splits into two steps: we first establish that, under the assumptions of Theorem 3, the error term

$$R_N(\alpha_{\text{ph}}, \Lambda) := \rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda) - \sum_{n=0}^N \nu_{n,\Lambda} \alpha_{\text{ph}}^n. \quad (5.3)$$

is controlled by a factor α_{ph}^{N+1} (up to some multiplicative constant depending only on N , ν and ϵ). In a second step we estimate the differences $\nu_{n,\Lambda} - \nu_n$ and deduce (12). More precisely, the remainder R_N satisfies the following

Lemma 5.1. *Let $N \in \mathbb{N}$ and $0 < \epsilon < 1$. Assume that $\epsilon \leq Z_3 = 1 - \alpha_{\text{ph}} B_\Lambda \leq 1 - \epsilon$ and $\mathcal{N}_N := \|(1+\mathcal{U})^{N+1}\nu\|_{L^2\cap\mathcal{C}} < +\infty$. Then, there exist two constants $C(m, \epsilon, \mathcal{N}_N)$ and $a(N, \epsilon, \mathcal{N}_N)$, depending only on N , ϵ and \mathcal{N}_N , such that*

$$\left\| R_N(\alpha_{\text{ph}}, \Lambda) \right\|_{L^2\cap\mathcal{C}} \leq C(N, \epsilon, \mathcal{N}_N) \alpha_{\text{ph}}^{N+1}, \quad (5.4)$$

for all $0 \leq \alpha_{\text{ph}} \leq a(N, \epsilon, \mathcal{N}_N)$.

As for the differences $\nu_{n,\Lambda} - \nu_n$, we have the

Lemma 5.2. *Let $\Lambda \geq 1$ and $N \in \mathbb{N}$. Assume that $\mathcal{N}_N := \|(1+\mathcal{U})^{N+1}\nu\|_{L^2\cap\mathcal{C}} < +\infty$. Then, there exists a constant $C(N, \mathcal{N}_N)$, depending only on N and \mathcal{N}_N , such that*

$$\left\| \nu_{n,\Lambda} - \nu_n \right\|_{L^2\cap\mathcal{C}} \leq \frac{C(N, \mathcal{N}_N)}{(1 + B_\Lambda)^{N+1-n}}, \quad (5.5)$$

for all $0 \leq n \leq N$.

Combining Lemmas 5.1 and 5.2, we can complete the proof of Theorem 3.

Proof of Theorem 3. Assumption (11), together with (A.3), means that

$$\mathcal{N}_N := \|(1 + \mathcal{U})^{N+1}\nu\|_{L^2 \cap \mathcal{C}} < +\infty.$$

It follows from (5.3) that

$$\rho_{\text{ph}}(\alpha_{\text{ph}}, \Lambda) - \sum_{n=0}^N \nu_n(\alpha_{\text{ph}})^n = R_N(\alpha_{\text{ph}}, \Lambda) + \sum_{n=0}^N (\nu_{n,\Lambda} - \nu_n)(\alpha_{\text{ph}})^n.$$

Hence by (5.4) and (5.5),

$$\left\| \rho_{\text{ph}}(\alpha_{\text{ph}}) - \sum_{n=0}^N \nu_n \alpha_{\text{ph}}^n \right\|_{L^2 \cap \mathcal{C}} \leq C(N, \epsilon, \mathcal{N}_N) \alpha_{\text{ph}}^{N+1} + C(N, \mathcal{N}_N) \sum_{n=0}^N \frac{\alpha_{\text{ph}}^n}{(1 + B_\Lambda)^{N+1-n}}, \quad (5.6)$$

for any number α_{ph} sufficiently small. In our setting we have $B_\Lambda \geq \epsilon/\alpha_{\text{ph}}$ and the result follows. \square

It therefore remains to show Lemmas 5.1 and 5.2.

Proof of Lemma 5.1. Let us introduce the notation

$$r_N(\alpha_{\text{ph}}) := (\alpha_{\text{ph}})^{-N-1} R_N(\alpha_{\text{ph}}). \quad (5.7)$$

We want to establish a bound on r_N independently of α_{ph} . By (5.3), this requires to estimate ρ_{ph} and $\nu_{n,\Lambda}$ (which was already done in Proposition 4.1).

The first step of the proof will be to bound ρ_{ph} independently of α_{ph} . Let us recall that a ground state for the reduced Bogoliubov-Dirac-Fock model satisfies

$$\|\alpha_{\text{ph}} \rho_{\text{ph}}\|_{\mathcal{C}} = \|\alpha(\rho_Q - \nu)\|_{\mathcal{C}} \leq \alpha \|\nu\|_{\mathcal{C}}$$

(see [16, Eq. (33)]). Since $\alpha_{\text{ph}} = Z_3 \alpha$, this provides

$$\|\rho_{\text{ph}}\|_{\mathcal{C}} \leq Z_3^{-1} \|\nu\|_{\mathcal{C}} \leq \epsilon^{-1} \|\nu\|_{\mathcal{C}}. \quad (5.8)$$

Note that we however do not have any a priori bound in $L^2(\mathbb{R}^3)$. Inserting (5.3) and (5.7) in (5.1) and using (2.8), we get

$$\begin{aligned} r_N &= \alpha_{\text{ph}} \mathcal{U}_\Lambda r_N + \mathcal{U}_\Lambda \nu_{N,\Lambda} + G_{N+1,\Lambda} + \sum_{k=N+1}^{N(N+2)} \alpha_{\text{ph}}^{k-N-1} \times \\ &\quad \times \sum_{3 \leq 2n+1 \leq N} \sum_{p_1 + \dots + p_{2n+1} = k-2n-1} F_{2n+1,\Lambda}(\omega_{p_1}, \dots, \omega_{p_{2n+1}}), \end{aligned} \quad (5.9)$$

where $\omega_p = \nu_{p,\Lambda}$ for $0 \leq p \leq N$, and $\omega_{N+1} = r_N$. It rests to estimate all the terms of the right-hand side of (5.9).

For the first term, we recall that $\alpha_{\text{ph}} |\mathcal{U}_\Lambda| \leq \alpha_{\text{ph}} B_\Lambda = 1 - Z_3 \leq 1 - \epsilon$, therefore

$$\|\alpha_{\text{ph}} \mathcal{U}_\Lambda r_N\|_{L^2 \cap \mathcal{C}} \leq (1 - \epsilon) \|r_N\|_{L^2 \cap \mathcal{C}}. \quad (5.10)$$

The second term can be controlled by using (3.3) and (4.1), which provide a positive constant $C(N, \mathcal{N}_N)$, depending only on N and \mathcal{N}_N , such that

$$\|\mathcal{U}_\Lambda \nu_{N,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq \kappa_1 \|(1 + \mathcal{U})\nu_{N,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq C(N, \mathcal{N}_N). \quad (5.11)$$

As for the function $G_{N+1,\Lambda}$, we first recall that

$$\left(1 - \frac{\pi^{\frac{1}{6}} 2^{\frac{11}{6}}}{\epsilon} \alpha_{\text{ph}} \|\nu\|_{\mathcal{C}}\right) |D^0| \leq \left|D^0 - \alpha_{\text{ph}} \rho_{\text{ph}} * \frac{1}{|\cdot|}\right| \leq \left(1 + \frac{\pi^{\frac{1}{6}} 2^{\frac{11}{6}}}{\epsilon} \alpha_{\text{ph}} \|\nu\|_{\mathcal{C}}\right) |D^0| \quad (5.12)$$

for all $\alpha_{\text{ph}} < \pi^{-1/6} 2^{-11/6} \epsilon \|\nu\|_{\mathcal{C}}^{-1}$ (see [16, p. 4495]). Hence, the operator $D^0 - \alpha_{\text{ph}} \rho_{\text{ph}} * |\cdot|^{-1}$ is invertible and, in particular, $G_{N+1,\Lambda}$ is well-defined. Notice also that (5.12) yields for any $\alpha_{\text{ph}} < \pi^{-1/6} 2^{-17/6} \epsilon \|\nu\|_{\mathcal{C}}^{-1}$

$$\frac{1}{2} |D^0| \leq \left|D^0 - \alpha_{\text{ph}} \rho_{\text{ph}} * \frac{1}{|\cdot|}\right| \leq \frac{3}{2} |D^0|.$$

When $N \geq 5$, we argue exactly as in Steps 1 and 2 of the proof of Proposition 3.2, and deduce that there exists a constant $C(N)$, depending only on N , such that

$$\|G_{N+1,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq C(N) \|\rho_{\text{ph}}\|_{\mathcal{C}}^{N+1}. \quad (5.13)$$

When $N \leq 4$, our argument is different. We expand $G_{N+1,\Lambda}$ as before, writing

$$G_{N+1,\Lambda} = -\alpha_{\text{ph}} \sum_{N+1 \leq 2j+1 \leq 5} F_{2j+1,\Lambda}(\rho_{\text{ph}}, \dots, \rho_{\text{ph}}) + G_{6,\Lambda}.$$

In view of (3.7) and (5.13) (for $N = 5$), this leads to

$$\|G_{N+1,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq C \alpha_{\text{ph}} \left(\sum_{N+1 \leq 2j+1 \leq 5} \|\rho_{\text{ph}}\|_{\mathcal{C}}^{2j+1} + \|\rho_{\text{ph}}\|_{\mathcal{C}}^6 \right).$$

In both cases, we obtain

$$\|G_{N+1,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq C(N) \max \left\{ \|\rho_{\text{ph}}\|_{\mathcal{C}}^{N+1}, \|\rho_{\text{ph}}\|_{\mathcal{C}}^6 \right\},$$

for any $\alpha_{\text{ph}} \leq 1$, so that, by (5.8),

$$\|G_{N+1,\Lambda}\|_{L^2 \cap \mathcal{C}} \leq C(N) \max \left\{ \frac{\|\nu\|_{\mathcal{C}}^{N+1}}{\epsilon^{N+1}}, \frac{\|\nu\|_{\mathcal{C}}^6}{\epsilon^6} \right\} \leq C(N, \mathcal{N}_N, \epsilon). \quad (5.14)$$

Finally, we consider the terms $\alpha_{\text{ph}}^{k-m-1} F_{2n+1,\Lambda}(\omega_{p_1}, \dots, \omega_{p_{2n+1}})$ of the sum in the right-hand side of (5.9). By (3.7), we have

$$\left\| \alpha_{\text{ph}}^{k-N-1} F_{2n+1,\Lambda}(\omega_{p_1}, \dots, \omega_{p_{2n+1}}) \right\|_{L^2 \cap \mathcal{C}} \leq C^{2n+1} |\alpha_{\text{ph}}|^{k-N-1} \prod_{j=1}^{2n+1} \|\omega_{p_j}\|_{\mathcal{C}}.$$

When $p_j \leq N$, we deduce from (4.1) that there exists a constant $C(N, \mathcal{N}_N)$ such that $\|\omega_{p_j}\|_{\mathcal{C}} \leq C(N, \mathcal{N}_N)$. On the other hand, when $p_j = N+1$ for some j , we can bound one of the norms $\|\omega_{p_j}\|_{\mathcal{C}}$ by $\|r_N\|_{\mathcal{C}}$, and the other ones by using (4.1), (5.3) and (5.8) to get

$$\|\omega_{p_j}\|_{\mathcal{C}} = |\alpha_{\text{ph}}|^{-p_j} \|R_N\| \leq C(N, \mathcal{N}_N, \epsilon) |\alpha_{\text{ph}}|^{-p_j},$$

for $\alpha_{\text{ph}} \leq 1$. This leads to

$$\left\| \alpha_{\text{ph}}^{k-N-1} F_{2n+1,\Lambda}(\omega_{p_1}, \dots, \omega_{p_{2n+1}}) \right\|_{L^2 \cap \mathcal{C}} \leq C(N, \mathcal{N}_N, \epsilon) |\alpha_{\text{ph}}|^{2n+1} \max\{\|r_N\|_{\mathcal{C}}, 1\},$$

for $\alpha_{\text{ph}} \leq 1$. Combining with (5.9), (5.10), (5.11) and (5.14), we conclude that

$$\|r_N\|_{L^2 \cap \mathcal{C}} \leq C(N, \mathcal{N}_N, \epsilon) + \left(1 - \epsilon + C(N, \mathcal{N}_N, \epsilon) |\alpha_{\text{ph}}|^3\right) \max\{\|r_N\|_{\mathcal{C}}, 1\},$$

for α_{ph} sufficiently small. Therefore, the norm $\|r_N\|_{L^2 \cap \mathcal{C}}$ is bounded independently of α_{ph} for α_{ph} small enough, which ends the proof of Lemma 5.1. \square

We finally prove Lemma 5.2.

Proof of Lemma 5.2. Given any $n \in \{0, 1, 2\}$, it follows from recursion relations (2.8) and (2.9) that

$$\nu_{n,\Lambda} - \nu_n = \mathcal{U}_\Lambda^n \nu_\Lambda - \mathcal{U}^n \nu = \mathcal{U}_\Lambda^n (\nu_\Lambda - \nu) + (\mathcal{U}_\Lambda^n - \mathcal{U}^n) \nu.$$

Therefore, given any $N \geq n$ and $0 \leq p \leq N + 1 - n$, we deduce from (3.3) that

$$\begin{aligned} \|(1 + \mathcal{U})^p (\nu_{n,\Lambda} - \nu_n)\|_{L^2 \cap \mathcal{C}} &\leq \kappa_1^n \|(1 + \mathcal{U})^{n+p} (\nu_\Lambda - \nu)\|_{L^2 \cap \mathcal{C}} \\ &\quad + n \kappa_1^{n-1} \|(1 + \mathcal{U})^{n+p-1} (\mathcal{U}_\Lambda - \mathcal{U}) \nu\|_{L^2 \cap \mathcal{C}}. \end{aligned} \quad (5.15)$$

Next, we recall that $\widehat{\nu}_\Lambda = \widehat{\nu} \mathbf{1}_{B(0, 2\Lambda)}$, so that, since $U(2\Lambda) = B_\Lambda$,

$$\|(1 + \mathcal{U})^{n+p} (\nu_\Lambda - \nu)\|_{L^2 \cap \mathcal{C}} \leq \frac{\|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}}}{(1 + U(2\Lambda))^{N+1-n-p}} = \frac{\|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}}}{(1 + B_\Lambda)^{N+1-n-p}}. \quad (5.16)$$

For the second term in the right-hand side of (5.15), we use (3.2) and write

$$\begin{aligned} \|(1 + \mathcal{U})^{n+p-1} (\mathcal{U}_\Lambda - \mathcal{U}) \nu\|_{L^2 \cap \mathcal{C}} &\leq \left\| \frac{U_\Lambda - U}{(1 + U)^{N+2-n-p}} \right\|_{L^\infty} \|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}} \\ &\leq \kappa_0^{N+4-n-p} \max \left\{ \frac{1}{(1 + B_\Lambda)^{N+1-n-p}}, \frac{1}{E(2\Lambda)} \right\} \|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}}. \end{aligned}$$

Since $(1 + B_\Lambda)^{N+1-n-p} \leq (1 + B_\Lambda)^{N+1} \leq C(N)E(2\Lambda)$, we obtain

$$\left\| (1 + \mathcal{U})^p (\nu_{n,\Lambda} - \nu_n) \right\|_{L^2 \cap \mathcal{C}} \leq \frac{C(N)}{(1 + B_\Lambda)^{N+1-n-p}} \|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}}.$$

Combining with (5.15) and (5.16), we are led to

$$\left\| (1 + \mathcal{U})^p (\nu_{n,\Lambda} - \nu_n) \right\|_{L^2 \cap \mathcal{C}} \leq \frac{C(N, \mathcal{N}_N)}{(1 + B_\Lambda)^{N+1-n-p}}, \quad (5.17)$$

for $N \geq n$ and $0 \leq p \leq N + 1 - n$.

We next turn to the case of $n \geq 3$. Given any $N \geq n$, we assume that (5.17) holds for all $n \leq k - 1$ and $0 \leq p \leq N + 1 - n$, and prove it by induction for $n = k$ and $0 \leq p \leq N + 1 - k$. Using (2.8) and (2.9), we first infer that

$$\begin{aligned} \|(1 + \mathcal{U})^p (\nu_{k,\Lambda} - \nu_k)\|_{L^2 \cap \mathcal{C}} &\leq \|(1 + \mathcal{U})^p \mathcal{U}_\Lambda (\nu_{k-1,\Lambda} - \nu_{k-1})\|_{L^2 \cap \mathcal{C}} \\ &\quad + \|(1 + \mathcal{U})^p (\mathcal{U}_\Lambda - \mathcal{U}) \nu_{k-1}\|_{L^2 \cap \mathcal{C}} + \sum_{3 \leq 2j+1 \leq k} \sum_{\substack{2j+1 \\ \ell=1}} \sum_{k_\ell = k-2j-1} \left\| (1 + \mathcal{U})^p \times \right. \\ &\quad \left. \times \left(F_{2j+1,\Lambda} (\nu_{k_1,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) - F_{2j+1} (\nu_{k_1}, \dots, \nu_{k_{2j+1}}) \right) \right\|_{L^2 \cap \mathcal{C}}. \end{aligned} \quad (5.18)$$

We next estimate the first term in the right-hand side of (5.18) using (3.3) and our assumption. This provides

$$\begin{aligned} \|(1 + \mathcal{U})^p \mathcal{U}_\Lambda (\nu_{k-1,\Lambda} - \nu_{k-1})\|_{L^2 \cap \mathcal{C}} &\leq \kappa_1 \|(1 + \mathcal{U})^{p+1} (\nu_{k-1,\Lambda} - \nu_{k-1})\|_{L^2 \cap \mathcal{C}} \\ &\leq \frac{C(N)}{(1 + B_\Lambda)^{N+1-k-p}} \|(1 + \mathcal{U})^{N+1} \nu\|_{L^2 \cap \mathcal{C}}. \end{aligned} \quad (5.19)$$

For the second term, we argue as in the proof of (5.17), using (3.2) and (2.12),

$$\begin{aligned} \|(1+U)^p(\mathcal{U}_\Lambda - U)\nu_{k-1}\|_{L^2\cap\mathcal{C}} &\leq \left\| \frac{U_\Lambda - U}{(1+U)^{N+2-k-p}} \right\|_{L^\infty} \|(1+U)^{N+2-k}\nu_{k-1}\|_{L^2\cap\mathcal{C}} \\ &\leq \frac{C(N, \mathcal{N}_N)}{(1+B_\Lambda)^{N+1-k-p}}. \end{aligned} \quad (5.20)$$

Finally, we turn to the terms in the sums of the right-hand side of (5.18). On the one hand, we deduce from (3.4) that

$$F_{2j+1,\Lambda} - F_{2j+1} = -F_{2j+1,\Lambda}^{(-1,1,\dots,1)} - F_{2j+1,\Lambda}^{(0,-1,1,\dots,1)} - \dots - F_{2j+1,\Lambda}^{(0,\dots,0,-1)}.$$

Hence, since $p \leq N+1-k \leq N+1-k_\ell$, we can apply (3.6) and (4.1) to obtain

$$\begin{aligned} \left\| (1+U)^p \left(F_{2j+1,\Lambda}(\nu_{k_1,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) - F_{2j+1}(\nu_{k_1,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) \right) \right\|_{L^2\cap\mathcal{C}} \\ \leq \frac{C(N)}{\Lambda^{\frac{1}{24}}} \prod_{\ell=1}^{2j+1} \|(1+U)^{N+1-k_\ell}\nu_{k_\ell,\Lambda}\|_{\mathcal{C}} \leq \frac{C(N, \mathcal{N}_N)}{\Lambda^{\frac{1}{24}}}. \end{aligned} \quad (5.21)$$

On the other hand, the multilinearity of the function F_{2j+1} provides

$$\begin{aligned} &F_{2j+1}(\nu_{k_1,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) - F_{2j+1}(\nu_{k_1}, \dots, \nu_{k_{2j+1}}) \\ &= F_{2j+1}(\nu_{k_1,\Lambda} - \nu_{k_1}, \nu_{k_2,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) + F_{2j+1}(\nu_{k_1}, \nu_{k_2,\Lambda} - \nu_{k_2}, \dots, \nu_{k_{2j+1},\Lambda}) \\ &\quad + \dots + F_{2j+1}(\nu_{k_1}, \nu_{k_2}, \dots, \nu_{k_{2j},\Lambda} - \nu_{k_{2j}}, \nu_{k_{2j+1},\Lambda}). \end{aligned}$$

Therefore, we infer similarly from (3.6) and (4.1) that

$$\begin{aligned} \left\| (1+U)^p \left(F_{2j+1}(\nu_{k_1,\Lambda}, \dots, \nu_{k_{2j+1},\Lambda}) - F_{2j+1}(\nu_{k_1}, \dots, \nu_{k_{2j+1}}) \right) \right\|_{L^2\cap\mathcal{C}} \\ \leq C(N) \sum_{q=1}^{2j+1} \left\| (1+U)^p(\nu_{k_q,\Lambda} - \nu_{k_q}) \right\|_{\mathcal{C}} \prod_{\ell < q} \left\| (1+U)^p \nu_{k_\ell,\Lambda} \right\|_{\mathcal{C}} \prod_{\ell > q} \left\| (1+U)^p \nu_{k_\ell} \right\|_{\mathcal{C}} \\ \leq \sum_{q=1}^{2j+1} \frac{C(N, \mathcal{N}_N)}{(1+B_\Lambda)^{N+1-k_q-p}} \leq \frac{C(N, \mathcal{N}_N)}{(1+B_\Lambda)^{N+1-k-p}}. \end{aligned} \quad (5.22)$$

As a conclusion, we derive from (5.18), (5.19), (5.20), (5.21) and (5.22) that

$$\left\| (1+U)^p(\nu_{k,\Lambda} - \nu_k) \right\|_{L^2\cap\mathcal{C}} \leq C(N, \mathcal{N}_N) \left(\frac{1}{(1+B_\Lambda)^{N+1-k-p}} + \frac{1}{\Lambda^{\frac{1}{24}}} \right),$$

Since $(1+B_\Lambda)^{N+1-k-p} \leq (1+B_\Lambda)^{N+1} \leq C(N)\Lambda^{1/24}$, this completes the proof of (5.17) for $n = k$.

Notice the constant $C(N, \mathcal{N}_N)$ deteriorates when N increases. However, this is not a problem since n is limited to the set $\{0, \dots, N\}$. Estimate (5.5) then follows from (5.17), considering the case $p = 0$. This concludes the proof of Lemma 5.2. \square

A Auxiliary results on the Uehling multiplier U

A.1 Elementary properties of U

We gather in this section some important properties of U , which will be useful for the proof of Lemma A.3 in the next section.

Lemma A.1. *The function U defined in (2.10) is a non-negative, non-decreasing, smooth function on \mathbb{R}_+ such that*

$$U(r) \underset{r \rightarrow 0}{\sim} \frac{r^2}{15\pi} \text{ and } U(r) \underset{r \rightarrow +\infty}{\sim} \frac{2}{3\pi} \log r. \quad (\text{A.1})$$

Its derivative U' is positive on $(0, +\infty)$, and it holds

$$U'(r) \underset{r \rightarrow +\infty}{\sim} \frac{2}{3\pi r} \text{ and } U''(r) \underset{r \rightarrow +\infty}{\sim} -\frac{2}{3\pi r^2}. \quad (\text{A.2})$$

Moreover, we have

$$\forall r \in \mathbb{R}^+, \frac{2}{15\pi}(1 + \log E(r)) \leq 1 + U(r) \leq 1 + \frac{2}{3\pi} \log E(r). \quad (\text{A.3})$$

Proof of Lemma A.1. For the convenience of the reader, let us recall the integral and the explicit formulas (2.10) of U

$$\begin{aligned} U(r) &= \frac{r^2}{4\pi} \int_0^1 \frac{z^2 - \frac{z^4}{3}}{1 + \frac{r^2(1-z^2)}{4}} dz \\ &= \frac{12 - 5r^2}{9\pi r^2} + \frac{\sqrt{4+r^2}}{3\pi r^3} (r^2 - 2) \log \left(\frac{\sqrt{4+r^2} + r}{\sqrt{4+r^2} - r} \right). \end{aligned} \quad (\text{A.4})$$

Most of the statements of Lemma A.1 are direct consequences of (A.4). As for (A.3), we estimate, using (A.4),

$$U(r) \leq \frac{r^2}{12\pi} \int_0^1 \frac{2z}{1 + \frac{r^2}{4} - \frac{r^2}{4} z^2} dz = \frac{1}{3\pi} \log \left(1 + \frac{r^2}{4} \right) \leq \frac{2}{3\pi} \log E(r).$$

For the lower bound, we notice similarly that

$$U(r) \geq \frac{r^2}{4\pi(1 + \frac{r^2}{4})} \int_0^1 \left(z^2 - \frac{z^4}{3} \right) dz = \frac{r^2}{15\pi(1 + \frac{r^2}{4})},$$

for $r \in \mathbb{R}_+$, so that

$$\forall 0 \leq r \leq 1, \frac{2}{15\pi}(1 + \log E(r)) \leq \frac{2+r^2}{15\pi} \leq 1 + U(r). \quad (\text{A.5})$$

On the other hand, we can also write

$$U(r) \geq \frac{r^2}{6\pi} \int_0^1 \frac{z^2}{1 + \frac{r^2}{2}(1-z)} dz = \frac{4}{3\pi r^4} \left(\left(1 + \frac{r^2}{2}\right)^2 \log \left(1 + \frac{r^2}{2}\right) - \frac{r^2}{2} - \frac{r^4}{8} \right),$$

thus when $r > 1$

$$1 + U(r) \geq \frac{1}{3\pi} \log \left(1 + \frac{r^2}{2} \right) + 1 - \frac{7}{6\pi} \geq \frac{1}{3\pi} (1 + \log E(r)).$$

The lower bound in (A.3) then follows from (A.5). \square

A useful consequence of Lemma A.1 is the following

Lemma A.2. Let Φ be the function defined on \mathbb{R}_+ by

$$\Phi(r) = \frac{U'(r)}{1 + U(r)},$$

There exist three positive numbers T_- , T_+ and Φ_0 such that the function Φ is an increasing diffeomorphism from $(0, T_-)$ onto $(0, \Phi_0)$, respectively a decreasing diffeomorphism from $(T_+, +\infty)$ onto $(0, \Phi_0)$, and $\Phi^{-1}((0, \Phi_0)) = (0, T_-) \cup (T_+, +\infty)$. Moreover, we have

$$\Phi(r) \underset{r \rightarrow 0}{\sim} \frac{2r}{15\pi}, \text{ and } \Phi(r) \underset{r \rightarrow +\infty}{\sim} \frac{1}{r \log r}. \quad (\text{A.6})$$

Proof of Lemma A.2. From Lemma A.1, we see that the function Φ is well-defined, smooth on \mathbb{R}_+ , and satisfies (A.6). Then we compute for $r \geq 0$:

$$\Phi'(r) = \frac{U''(r)(1 + U(r)) - U'(r)^2}{(1 + U(r))^2}.$$

By (A.1) and (A.2), we thus have $\Phi'(0) = \frac{2}{15\pi}$ and $\Phi'(r) \underset{r \rightarrow +\infty}{\sim} -1/(r^2 \log r)$. Since $\Phi(0) = 0$ and $\Phi(r) \rightarrow 0$ as $r \rightarrow +\infty$ by (A.1) and (A.2), there exist $a, b, \delta > 0$ such that Φ is an increasing diffeomorphism from $(0, a)$ onto $(0, \delta)$, respectively a decreasing diffeomorphism from $(b, +\infty)$ onto $(0, \delta)$. The function Φ is positive on $[a, b]$, so that $m = \min\{\Phi(t), a \leq t \leq b\} > 0$. Lemma A.2 follows by introducing $\Phi_0 = \min\{m/2, \delta\}$, and $T_- < T_+$, the two positive numbers such that $\Phi(T_-) = \Phi(T_+) = \Phi_0$. \square

A.2 A useful bound involving U

We use here results from the previous section to derive a bound useful for the proof of Proposition 3.2.

Lemma A.3. *There exists a universal constant $K > 0$ such that*

$$1 + U \left(\left| \sum_{j=1}^n v_j \right| \right) \leq K \log n \prod_{j=1}^n (1 + U(|v_j|)) \quad (\text{A.7})$$

for all $n \geq 1$, and all $(v_1, \dots, v_n) \in (\mathbb{R}^3)^n$.

If we allow K to depend on n , the optimal constant in the above inequality satisfies $K_n \rightarrow 1/3\pi$ when $n \rightarrow +\infty$, as can be seen from the proof. The factor $\log n$ in (A.7) is therefore optimal with regard to the large- n dependence.

Proof of Lemma A.3. By Lemma A.1, we have

$$\frac{1 + U \left(\left| \sum_{j=1}^n v_j \right| \right)}{\prod_{j=1}^n (1 + U(|v_j|))} \leq \frac{1 + U \left(\sum_{j=1}^n |v_j| \right)}{\prod_{j=1}^n (1 + U(|v_j|))} \leq \max_{(t_1, \dots, t_n) \in \mathbb{R}_+^n} \frac{1 + U \left(\sum_{j=1}^n t_j \right)}{\prod_{j=1}^n (1 + U(t_j))} := J_n. \quad (\text{A.8})$$

It is clear that taking $v_1 = \dots = v_n = v$ shows that the maximum of the left-hand side of (A.8) is actually J_n . Next, we take $t_1 = \dots = t_n = \tau_n$ in (A.8) with $\tau_n = \sqrt{15\pi/(n \log n)}$. Using (A.2), we see that $J_n \gtrsim (\log n)/3\pi$ for $n \gg 1$. We will show that actually it holds $J_n \sim (\log n)/(3\pi)$ when $n \rightarrow +\infty$. In the rest of the proof, we assume $n \geq n_0$ is such that $J_n > 1$.

Let us consider a maximizing sequence $\{(t_1^{(p)}, \dots, t_n^{(p)})\}_{p \in \mathbb{N}}$ for the variational problem defining J_n . If the sequence is unbounded, then by Lemma A.1,

$$J_n \leq \lim_{p \rightarrow +\infty} \frac{1 + U\left(n \max_j \{t_j^{(p)}\}\right)}{1 + U\left(\max_j \{t_j^{(p)}\}\right)} = 1$$

which contradicts $J_n > 1$ for $n \geq n_0$. Therefore $(t_1^{(p)}, \dots, t_n^{(p)})$ is bounded in $(\mathbb{R}_+)^n$. In this case the variational problem on the right-hand side of (A.8) has a maximizer, which satisfies the equation

$$\forall 1 \leq k \leq n, \Phi(t_k) = \frac{U'(t_k)}{1 + U(t_k)} = \frac{U'\left(\sum_{j=1}^n t_j\right)}{1 + U\left(\sum_{j=1}^n t_j\right)} = \Phi\left(\sum_{j=1}^n t_j\right) := \Phi_1. \quad (\text{A.9})$$

Assume now that $\Phi_1 \geq \Phi_0$. By Lemma A.2, we have $t_k \geq T_-$ and $\sum_{j=1}^n t_j \leq T_+$, for all $1 \leq k \leq n$, hence $n \leq T_+/T_-$. In particular, for $n > T_+/T_-$, it must hold $0 \leq \Phi_1 < \Phi_0$. If $\Phi_1 = 0$, we infer from Lemma A.1 that $t_1 = \dots = t_n = 0$, so that $J_n = 1$, a contradiction. Therefore, by Lemma A.2, there exist exactly two numbers $0 < \tau_n < T_-$ and $T_n > T_+$ such that $\Phi(\tau_n) = \Phi(T_n) = \Phi_1$. By (A.9), the unique possible maximizer is (τ_n, \dots, τ_n) , where $\tau_n = T_n/n \in (0, T_-)$ is such that

$$\Phi(\tau_n) = \Phi(n\tau_n). \quad (\text{A.10})$$

The corresponding value of J_n is

$$J_n = \frac{1 + U(n\tau_n)}{(1 + U(\tau_n))^n}. \quad (\text{A.11})$$

By (A.10), we must have $\tau_n \rightarrow 0$ as $n \rightarrow +\infty$. Combining (A.10) with (A.6), it follows that $\Phi(n\tau_n) \sim 2\tau_n/15\pi \rightarrow 0$. By Lemma A.2 and since $n\tau_n = T_n \geq T_+$, it holds $n\tau_n \rightarrow +\infty$. Using (A.6) again, we deduce that $n\tau_n \log(n\tau_n) \sim_{n \rightarrow +\infty} 15\pi/2\tau_n$, hence finally $\tau_n \sim \sqrt{15\pi/(n \log n)}$. Inserting in (A.1) and (A.11), we finally arrive at $J_n \sim (\log n)/(3\pi)$. This ends the proof of Lemma A.3. \square

B Proof of Proposition 3.1

We start by showing the following lemma which provides estimates on $U - U_\Lambda$ in $[0, 2\Lambda]$.

Lemma B.1. *Let $\Lambda \geq 1$. For $\kappa_1 = 258/\pi$, one has*

$$\forall 0 \leq r \leq 2\Lambda, |U_\Lambda(r) - U(r)| \leq \kappa_1 \frac{r}{2E(\Lambda)}. \quad (\text{B.1})$$

Proof of Lemma B.1. Recall that (see (2.10) and (3.1))

$$\begin{aligned} U_\Lambda(r) - U(r) &= \frac{r^2}{4\pi} \int_{\frac{\Lambda}{E(\Lambda)}}^1 \frac{z^2 - \frac{z^4}{3}}{1 + \frac{r^2}{4}(1 - z^2)} dz - \frac{r}{2\pi} \int_0^{Z_\Lambda(r)} \frac{z - \frac{z^3}{3}}{E(\Lambda) - \frac{rz}{2}} dz \\ &\quad + \frac{1}{\pi} \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{(1 - z^2)(1 + \frac{r^2}{4}(1 - z^2))} dz, \end{aligned} \quad (\text{B.2})$$

for $0 \leq r \leq 2\Lambda$ and where

$$Z_\Lambda(r) = \frac{E(\Lambda) - E(\Lambda - r)}{r} = \frac{2\Lambda - r}{E(\Lambda) + E(\Lambda - r)} \leq \frac{\Lambda}{E(\Lambda)}. \quad (\text{B.3})$$

We will estimate all the terms of the right-hand side of (B.2). The first term is treated as follows, for all $0 \leq r \leq 2\Lambda$:

$$\left| \frac{r^2}{4\pi} \int_{\frac{\Lambda}{E(\Lambda)}}^1 \frac{z^2 - \frac{z^4}{3}}{1 + \frac{r^2}{6}(1 - z^2)} dz \right| \leq \frac{r^2}{6\pi} \left(1 - \frac{\Lambda}{E(\Lambda)}\right) \leq \frac{r^2}{6\pi E(\Lambda)^2}. \quad (\text{B.4})$$

Using (B.3) and $|x| \leq E(x)$, we bound the second term by

$$\frac{r}{2\pi} \left| \int_0^{Z_\Lambda(r)} \frac{z - \frac{z^3}{3}}{E(\Lambda) - \frac{rz}{2}} dz \right| \leq \frac{rZ_\Lambda(r)^2}{4\pi(E(\Lambda) - \frac{rZ_\Lambda(r)}{2})} \leq \frac{r}{2\pi E(\Lambda)}, \quad (\text{B.5})$$

for $0 \leq r \leq 2\Lambda$. In order to estimate the last term of the right-hand side of (B.2), we distinguish the regions $0 \leq r \leq \Lambda/2$ and $\Lambda/2 \leq r \leq 2\Lambda$. We calculate

$$\left| \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{(1 - z^2)(1 + \frac{r^2}{4}(1 - z^2))} dz \right| \leq \frac{2}{3} \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{dz}{1 - z} = \frac{2}{3} \log \left(\frac{1 - Z_\Lambda(r)}{1 - \frac{\Lambda}{E(\Lambda)}} \right).$$

On the other hand, by (B.3),

$$\frac{1 - Z_\Lambda(r)}{1 - \frac{\Lambda}{E(\Lambda)}} = 1 + \frac{r(2\Lambda - r)(\Lambda + E(\Lambda))}{(E(\Lambda) + E(\Lambda - r))((\Lambda - r)E(\Lambda) + \Lambda E(\Lambda - r))} \leq 1 + \frac{6r}{E(\Lambda)},$$

as soon as $0 \leq r \leq \Lambda/2$. Hence using $\log(1 + x) \leq x$ we infer the bound

$$\forall 0 \leq r \leq \frac{\Lambda}{2}, \frac{1}{\pi} \left| \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{(1 - z^2)(1 + \frac{r^2}{4}(1 - z^2))} dz \right| \leq \frac{4r}{\pi E(\Lambda)}. \quad (\text{B.6})$$

For $\Lambda/2 \leq r \leq 2\Lambda$, we write similarly as before

$$\left| \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{(1 - z^2)(1 + \frac{r^2}{4}(1 - z^2))} dz \right| \leq \frac{8}{3r^2} \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{dz}{(1 - z)^2} \leq \frac{8}{3r^2} E(\Lambda)(\Lambda + E(\Lambda))$$

and deduce the estimate

$$\forall \frac{\Lambda}{2} \leq r \leq 2\Lambda, \frac{1}{\pi} \left| \int_{Z_\Lambda(r)}^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{(1 - z^2)(1 + \frac{r^2}{4}(1 - z^2))} dz \right| \leq \frac{128r}{\pi E(\Lambda)}. \quad (\text{B.7})$$

Estimate (B.1) follows from (B.4), (B.5), (B.6) and (B.7), together with (B.2). This ends the proof of Lemma B.1. \square

We now use Lemma B.1 to finish the proof of Proposition 3.1. The pointwise convergence of U_Λ when $\Lambda \rightarrow +\infty$ is a direct consequence of (B.1). For (3.2), we first use (A.3) and (B.1) to obtain

$$\forall 0 \leq r \leq 2\Lambda, \left| \frac{U_\Lambda(r) - U(r)}{(1 + U(r))^{m+1}} \right| \leq \kappa_1 \left(\frac{15\pi}{2} \right)^{m+1} \frac{E(r)}{2E(\Lambda)(1 + \log E(r))^{m+1}}.$$

Optimizing $x \mapsto \frac{E(x)}{(1 + \log E(x))^{m+1}}$ on $[0, 2\Lambda]$ yields

$$\frac{E(r)}{(1 + \log E(r))^{m+1}} \leq \max \left\{ 1, \frac{E(2\Lambda)}{(1 + \log E(2\Lambda))^{m+1}} \right\}.$$

Since $E(2x) \leq 2E(x)$ for any $x \geq 0$, we are led to

$$\left| \frac{U_\Lambda(r) - U(r)}{(1 + U(r))^{m+1}} \right| \leq \kappa_1 \left(\frac{15\pi}{2} \right)^{m+1} \max \left\{ \frac{1}{(1 + \log E(2\Lambda))^{m+1}}, \frac{1}{E(2\Lambda)} \right\}. \quad (\text{B.8})$$

On the other hand, U is non-decreasing on \mathbb{R}_+ , hence, using (A.3) we infer

$$\forall r \geq 2\Lambda, \left| \frac{U_\Lambda(r) - U(r)}{(1 + U(r))^{m+1}} \right| = \left| \frac{U(r)}{(1 + U(r))^{m+1}} \right| \leq \left(\frac{15\pi}{2} \right)^m \frac{1}{(1 + \log E(2\Lambda))^m}.$$

Using (B.8), we finally obtain

$$\left\| \frac{U_\Lambda - U}{(1 + U)^{m+1}} \right\|_{L^\infty} \leq \kappa_1 \left(\frac{15\pi}{2} \right)^{m+1} \max \left\{ \frac{1}{(1 + \log E(2\Lambda))^m}, \frac{1}{E(2\Lambda)} \right\}. \quad (\text{B.9})$$

We now recall that

$$B_\Lambda = \frac{1}{\pi} \int_0^{\frac{\Lambda}{E(\Lambda)}} \frac{z^2 - \frac{z^4}{3}}{1 - z^2} dz, \quad (\text{B.10})$$

so that, for $\Lambda \geq 1$,

$$B_\Lambda \leq \frac{2}{3\pi} \int_0^{\frac{\Lambda}{E(\Lambda)}} \frac{dz}{1 - z} = \frac{2}{3\pi} \log [E(\Lambda)(\Lambda + E(\Lambda))] \leq \frac{4}{3\pi} \log E(2\Lambda).$$

Combining with (B.9), we finally derive (3.2). We end the proof of Proposition 3.1 by noting that (3.3) follows directly from the definition of U_Λ and (B.1).

C Proof of Proposition 3.2

We may define $F_{n,\Lambda}^\epsilon(\mu)$ by duality as follows

$$\int_{\mathbb{R}^3} \zeta F_{n,\Lambda}^\epsilon(\mu) = \text{tr}(Q_{n,\Lambda}^\epsilon \zeta),$$

for any smooth function ζ , and where

$$Q_{n,\Lambda}^\epsilon = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{D^0 + i\eta} \prod_{j=1}^n \left(\Pi_\Lambda^{(\epsilon_j)} \mu_j * \frac{1}{|x|} \Pi_\Lambda^{(\epsilon_{j+1})} \frac{1}{D^0 + i\eta} \right) d\eta.$$

We will use, like in [15, p. 547], the inequality

$$|\text{tr}(Q_{n,\Lambda}^\epsilon \zeta)| = \left| \int_{\mathbb{R}^3} \text{tr}_{\mathbb{C}^4} \left(\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p) \right) dp \right| \leq \int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp. \quad (\text{C.1})$$

The main idea is to derive a bound of the last integral in (C.1) in terms of the norms $\|(1 + \mathcal{U})^{-m} \zeta\|_L^2$ and $\|(1 + \mathcal{U})^{-m} \zeta\|_{L^2 + \mathcal{C}'}$, which provides an estimate of the form (3.6), by duality. The proof will depend on whether we estimate the integral in the right-hand side of (C.1) by the norm $\|(1 + \mathcal{U})^{-m} \zeta\|_{\mathcal{C}'}$ or by the norm $\|(1 + \mathcal{U})^{-m} \zeta\|_{L^2}$. For this reason, we split it into three steps.

Step 1. *There exists a universal constant C_1 such that for all $n \geq 5$*

$$\|(1 + \mathcal{U})^m F_{n,\Lambda}^\epsilon(\mu)\|_{\mathcal{C}} \leq \frac{(C_1)^n (K \log n)^m}{\Lambda^{\frac{n(\epsilon)}{2}}} \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (\text{C.2})$$

for all $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{C}^n$.

We estimate $\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)$ as follows:

$$\begin{aligned}
|\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| &\leq \frac{1}{(2\pi)^{\frac{3n+5}{2}}} \int_{-\infty}^{+\infty} d\eta \int_{\mathbb{R}^3} \cdots \int_{\mathbb{R}^3} \left(f_{\frac{1}{4}}^{(\epsilon_1)}(p) |\widehat{\phi}_1(p-p_1)| f_{\frac{1}{4}}^{(\epsilon_2)}(p_1) \right) \times \\
&\times \prod_{j=1}^{n-1} \left(f_{\frac{1}{4}}^{(\epsilon_{j+1})}(p_j) |\widehat{\phi}_{j+1}(p_j-p_{j+1})| f_{\frac{1}{4}}^{(\epsilon_{j+2})}(p_{j+1}) \right) \times \\
&\times \left(f_{\frac{1}{4}}^{(\epsilon_{n+1})}(p_n) |\widehat{\zeta}(p_n-p)| f_{\frac{1}{4}}^{(\epsilon_1)}(p) \right) dp_1 \cdots dp_n,
\end{aligned} \tag{C.3}$$

where $\phi_j = \mu_j * |\cdot|^{-1}$, and for any $\beta > 0$, $f_\beta^{(\epsilon)} = \pi_\Lambda^{(\epsilon)} / (\eta^2 + E^2)^\beta$, with $\pi_\Lambda^{(\epsilon)} = 1$, if $\epsilon \neq -1$, and $\pi_\Lambda^{(-1)}(\cdot) = \mathbb{1}_{|\cdot| > \Lambda}$. Applying the following corollary of (A.7)

$$(1 + U(p - p_n))^m \leq (K \log n)^m (1 + U(p - p_1))^m \prod_{j=1}^{n-1} (1 + U(p_j - p_{j+1}))^m$$

to (C.3), we are led to estimating

$$\begin{aligned}
\int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp &\leq \frac{1}{2\pi} (K \log n)^m \int_{-\infty}^{+\infty} \text{tr} \left(\prod_{j=1}^n (f_{\frac{1}{4}}^{(\epsilon_j)}(-i\nabla) \psi_j(x) \times \right. \\
&\times \left. f_{\frac{1}{4}}^{(\epsilon_{j+1})}(-i\nabla) (f_{\frac{1}{4}}^{(\epsilon_{n+1})}(-i\nabla) \xi(x) f_{\frac{1}{4}}^{(\epsilon_1)}(-i\nabla)) \right) d\eta,
\end{aligned}$$

where $\widehat{\psi}_j = (1 + U)^m |\widehat{\phi}_j|$ for $1 \leq j \leq n$, and $\widehat{\xi} = (1 + U)^{-m} |\widehat{\zeta}|$. Since $n + 1 \geq 6$, we deduce from Hölder's inequality in Schatten spaces [30], and the fact that $\|\cdot\|_{\mathfrak{S}_q} \leq \|\cdot\|_{\mathfrak{S}_r}$, as soon as $1 \leq r \leq q \leq +\infty$, that

$$\begin{aligned}
\int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp &\leq \frac{1}{2\pi} (K \log n)^m \int_{-\infty}^{+\infty} \left(\prod_{j=1}^n \|f_{\frac{1}{4}}^{(\epsilon_j)}(-i\nabla) \psi_j(x) \times \right. \\
&\times \left. f_{\frac{1}{4}}^{(\epsilon_{j+1})}(-i\nabla) \right\|_{\mathfrak{S}_6} \|f_{\frac{1}{4}}^{(\epsilon_{n+1})}(-i\nabla) \xi(x) f_{\frac{1}{4}}^{(\epsilon_1)}(-i\nabla)\|_{\mathfrak{S}_6} \right) d\eta.
\end{aligned} \tag{C.4}$$

We now use the Kato-Seiler-Simon inequality (see [27] and [30, Thm 4.1]),

$$\forall p \geq 2, \|f(-i\nabla)g(x)\|_{\mathfrak{S}_p} \leq \frac{1}{(2\pi)^{\frac{p}{3}}} \|g\|_{L^p(\mathbb{R}^3)} \|f\|_{L^p(\mathbb{R}^3)}, \tag{C.5}$$

to bound all the terms of the product in the right-hand side of (C.4). This provides

$$\begin{aligned}
\|f_{\frac{1}{4}}^{(\epsilon)}(-i\nabla) |h(x)| f_{\frac{1}{4}}^{(\epsilon')}(-i\nabla)\|_{\mathfrak{S}_6} &\leq \|f_{\frac{1}{4}}^{(\epsilon)}(-i\nabla) |h(x)|^{\frac{1}{2}}\|_{\mathfrak{S}_{12}} \| |h(x)|^{\frac{1}{2}} f_{\frac{1}{4}}^{(\epsilon')}(-i\nabla)\|_{\mathfrak{S}_{12}} \\
&\leq \frac{1}{(2\pi)^{\frac{1}{2}}} \|h\|_{L^6} \|f_{\frac{1}{4}}^{(\epsilon)}\|_{L^{12}} \|f_{\frac{1}{4}}^{(\epsilon')}\|_{L^{12}},
\end{aligned}$$

for any ϵ and ϵ' in $\{-1, 0, 1\}$, and any $h \in L^6(\mathbb{R}^3)$. In particular, by the critical Sobolev inequality, we obtain for any function h in $\dot{H}^1(\mathbb{R}^3)$,

$$\|f_{\frac{1}{4}}^{(\epsilon)}(-i\nabla) |h(x)| f_{\frac{1}{4}}^{(\epsilon')}(-i\nabla)\|_{\mathfrak{S}_6} \leq A \|\nabla h\|_{L^2} \|f_{\frac{1}{4}}^{(\epsilon)}\|_{L^{12}} \|f_{\frac{1}{4}}^{(\epsilon')}\|_{L^{12}}, \tag{C.6}$$

for some universal constant A . Given any $q \geq 2$ and $\beta > 6/q$, we then check that

$$\begin{aligned}
\|f_\beta^{(\epsilon)}\|_{L^q} &\leq E(\eta)^{\frac{3}{q}-2\beta} \left(\int_{\mathbb{R}^3} \frac{du}{E(u)^{\beta q}} \right)^{\frac{1}{q}} \leq E(\eta)^{\frac{3}{q}-2\beta} \left(\int_{|u| \geq 1} \frac{du}{|u|^{2\beta q}} \right)^{\frac{1}{q}} \\
&= \left(\frac{4\pi}{2\beta q - 3} \right)^{\frac{1}{q}} E(\eta)^{\frac{3}{q}-2\beta},
\end{aligned} \tag{C.7}$$

for $\epsilon \neq -1$, while similarly,

$$\|f_\beta^{(-1)}\|_{L^q} \leq \left(\frac{4\pi}{2\beta q - 3}\right)^{\frac{1}{q}} \min \left\{ E(\eta)^{\frac{3}{q} - 2\beta}, \Lambda^{\frac{3}{q} - 2\beta} \right\}. \quad (\text{C.8})$$

The definition of the functions ψ_j gives

$$\|(1 + \mathcal{U})^m \nabla \psi_j\|_{L^2} = 4\pi \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (\text{C.9})$$

which, combined with (C.4), (C.6), (C.7) and (C.8), leads to

$$\begin{aligned} \int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp &\leq A^{n+1} (K \log n)^m \|(1 + \mathcal{U})^{-m} \zeta\|_{\mathcal{C}'} \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}} \times \\ &\quad \times \int_0^{+\infty} \min \left\{ \frac{1}{E(\eta)^{\frac{1}{2}}}, \frac{1}{\Lambda^{\frac{1}{2}}} \right\}^{n_\epsilon} \frac{d\eta}{E(\eta)^{\frac{n+1-n_\epsilon}{2}}}, \end{aligned}$$

for some universal constant A . When $n_\epsilon = 0$, we have $\int_0^{+\infty} E(\eta)^{-(n+1)/2} d\eta \leq \int_0^{+\infty} E(\eta)^{-3} d\eta$, whereas, for $n_\epsilon = 1$,

$$\int_0^{+\infty} \min \left\{ \frac{1}{E(\eta)^{\frac{1}{2}}}, \frac{1}{\Lambda^{\frac{1}{2}}} \right\} \frac{d\eta}{E(\eta)^{\frac{n}{2}}} \leq \frac{1}{\Lambda^{\frac{1}{2}}} \int_0^{+\infty} \frac{d\eta}{E(\eta)^{\frac{5}{2}}} + \frac{1}{2\Lambda^2}.$$

Inequality (C.2) then follows with $C_1 = A/2 + A \int_0^{+\infty} E(\eta)^{-5/2} d\eta$.

Step 2. *There exists a universal constant C_2 such that, for $n = 3$ or $n \geq 5$,*

$$\|(1 + \mathcal{U})^m F_{n,\Lambda}^\epsilon(\mu)\|_{L^2(\mathbb{R}^3)} \leq \frac{(C_2)^n (K \log n)^m}{\Lambda^{\frac{n(\epsilon)}{7}}} \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (\text{C.10})$$

for all $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{C}^n$.

The proof is similar to the proof of (C.2). Since $n/6 + 1/2 \geq 1$, we can now estimate $\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)$ by

$$\begin{aligned} \int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp &\leq \frac{(K \log n)^m}{2\pi} \int_{-\infty}^{+\infty} \left(\|\psi_1(x) f_{\frac{3}{8}}^{(\epsilon_2)}(-i\nabla)\|_{\mathfrak{S}_6} \times \right. \\ &\quad \times \|f_{\frac{1}{8}}^{(\epsilon_2)}(-i\nabla) \psi_2(x) f_{\frac{1}{4}}^{(\epsilon_3)}(-i\nabla)\|_{\mathfrak{S}_6} \prod_{j=2}^{n-2} \|f_{\frac{1}{4}}^{(\epsilon_{j+1})}(-i\nabla) \psi_{j+1}(x) f_{\frac{1}{4}}^{(\epsilon_{j+2})}(-i\nabla)\|_{\mathfrak{S}_6} \\ &\quad \left. \times \|f_{\frac{1}{4}}^{(\epsilon_n)}(-i\nabla) \psi_n(x) f_{\frac{1}{8}}^{(\epsilon_{n+1})}(-i\nabla)\|_{\mathfrak{S}_6} \|f_{\frac{3}{8}}^{(\epsilon_{n+1})}(-i\nabla) \xi(x) f_{\frac{1}{2}}^{(\epsilon_1)}(-i\nabla)\|_{\mathfrak{S}_2} \right) d\eta, \end{aligned} \quad (\text{C.11})$$

where the functions $f_\beta^{(\epsilon)}$, ψ_j and ξ are defined as in Step 1. Using Hölder's inequality and (C.5), we can bound each norm in the right-hand side of (C.11) similarly to (C.6). This provides, for instance,

$$\|f_{\frac{1}{8}}^{(\epsilon_2)}(-i\nabla) |h(x)| f_{\frac{1}{4}}^{(\epsilon_3)}(-i\nabla)\|_{\mathfrak{S}_6} \leq A \|\nabla h\|_{L^2} \|f_{\frac{1}{8}}^{(\epsilon_2)}\|_{L^{18}} \|f_{\frac{1}{4}}^{(\epsilon_3)}\|_{L^9},$$

and

$$\|f_{\frac{3}{8}}^{(\epsilon_{n+1})}(-i\nabla) |h(x)| f_{\frac{1}{2}}^{(\epsilon_1)}(-i\nabla)\|_{\mathfrak{S}_2} \leq \frac{1}{(2\pi)^{\frac{3}{2}}} \|h\|_{L^2} \|f_{\frac{3}{8}}^{(\epsilon_{n+1})}\|_{L^{\frac{14}{3}}} \|f_{\frac{1}{2}}^{(\epsilon_1)}\|_{L^{\frac{7}{2}}}.$$

Combining with (C.7), (C.8) and (C.9), we obtain

$$\int_{\mathbb{R}^3} |\widehat{Q_{n,\Lambda}^\epsilon \zeta}(p, p)| dp \leq A^{n+1} (K \log n)^m \|(1 + \mathcal{U})^{-m} \zeta\|_{L^2} \prod_{j=1}^n \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}} \times \\ \times \int_0^{+\infty} \min \left\{ \frac{1}{E(\eta)}, \frac{1}{\Lambda} \right\}^{\frac{n_\epsilon}{7}} \frac{d\eta}{E(\eta)^{\frac{3n-2}{6} + \frac{1-n_\epsilon}{7}}},$$

for a universal constant A . Since

$$\int_0^{+\infty} \frac{d\eta}{E(\eta)^{\frac{3n-2}{6} + \frac{1}{7}}} \leq \int_0^{+\infty} \frac{d\eta}{E(\eta)^{\frac{7}{6}}},$$

for $n_\epsilon = 0$, whereas, for $n_\epsilon = 1$,

$$\int_0^{+\infty} \min \left\{ \frac{1}{E(\eta)^{\frac{1}{7}}}, \frac{1}{\Lambda^{\frac{1}{7}}} \right\} \frac{d\eta}{E(\eta)^{\frac{3n-2}{6}}} \leq \frac{1}{\Lambda^{\frac{1}{7}}} \int_0^{+\infty} \frac{d\eta}{E(\eta)^{\frac{7}{6}}} + \frac{6}{\Lambda^{\frac{1}{6}}},$$

we obtain (C.10) with $C_2 = 6A + A \int_0^{+\infty} E(\eta)^{-7/6} d\eta$.

Step 3. Let $n = 3$. There exists a universal constant C_3 such that

$$\|(1 + \mathcal{U})^m F_{3,\Lambda}^\epsilon(\mu)\|_{\mathcal{C}} \leq \frac{(C_3)^3 (K \log 3)^m}{\Lambda^{\frac{n_\epsilon}{24}}} \prod_{j=1}^3 \|(1 + \mathcal{U})^m \mu_j\|_{\mathcal{C}}, \quad (\text{C.12})$$

for all $\mu = (\mu_1, \mu_2, \mu_3) \in \mathcal{C}^3$.

The proof of (C.12) follows ideas from [15, Section 4.3.4]. Contrarily to Steps 1 and 2, it relies on an explicit computation of $\widehat{Q_{3,\Lambda}^\epsilon \zeta}(p, p)$ by means of the residuum formula for the integral with respect to the variable η . Indeed, it holds

$$\widehat{Q_{3,\Lambda}^\epsilon \zeta}(p, p) = \sum_{\delta \in \{-1, 1\}^4} \widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p). \quad (\text{C.13})$$

Here, the quantity $\widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p)$ vanishes if $\delta = \pm(1, 1, 1, 1)$, whereas, when $\delta = (1, -1, -1, -1)$, it refers to the expression

$$\widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p) = \frac{1}{(2\pi)^6} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \pi_\Lambda^{(\epsilon_1)}(p) P_+^0(p) \frac{\widehat{\phi}_1(p - p_1)}{E(p) + E(p_1)} \pi_\Lambda^{(\epsilon_2)}(p_1) P_-^0(p_1) \times \\ \times \frac{\widehat{\phi}_2(p_1 - p_2)}{E(p) + E(p_2)} \pi_\Lambda^{(\epsilon_3)}(p_2) P_-^0(p_2) \frac{\widehat{\phi}_3(p_2 - p_3)}{E(p) + E(p_3)} \pi_\Lambda^{(\epsilon_4)}(p_3) P_-^0(p_3) \widehat{\zeta}(p_3 - p) dp_1 dp_2 dp_3,$$

where $P_\pm^0(p) = (E(p) \pm (\alpha \cdot p + \beta)) / 2E(p)$. The expression of $\widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p)$ is similar when δ contains exactly one $\delta_i = 1$, respectively exactly one $\delta_i = -1$. On the other hand, for $\delta = (1, 1, -1, -1)$, the function $\widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p)$ is given by

$$\widehat{Q_{3,\Lambda}^{\epsilon, \delta} \zeta}(p, p) = \frac{1}{(2\pi)^6} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \pi_\Lambda^{(\epsilon_1)}(p) P_+^0(p) |\widehat{\phi}_1(p - p_1)| \pi_\Lambda^{(\epsilon_2)}(p_1) P_-^0(p_1) \times \\ \times |\widehat{\phi}_2(p_1 - p_2)| \pi_\Lambda^{(\epsilon_3)}(p_2) P_-^0(p_2) |\widehat{\phi}_3(p_2 - p_3)| \pi_\Lambda^{(\epsilon_4)}(p_3) P_-^0(p_3) \widehat{\zeta}(p_3 - p) \times \\ \times \left(\frac{1}{(E(p) + E(p_2))(E(p_1) + E(p_2))(E(p_1) + E(p_3))} \right. \\ \left. + \frac{1}{(E(p) + E(p_2))(E(p) + E(p_3))(E(p_1) + E(p_3))} \right) dp_1 dp_2 dp_3.$$

We next estimate $\widehat{Q_{3,\Lambda}^{\epsilon,\delta}\zeta}(p, p)$ as above. For instance, when $\delta = (1, -1, -1, -1)$, since $E(p + q) \leq E(p) + E(q)$ for any $(p, q) \in \mathbb{R}^3$, we can compute

$$\begin{aligned} |\widehat{Q_{3,\Lambda}^{\epsilon,\delta}\zeta}(p, p)| &\leq \frac{1}{(2\pi)^6} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} dp_1 dp_2 dp_3 \frac{|P_+^0(p)\widehat{\phi}_1(p-p_1)P_-^0(p_1)|}{E(p+p_1)^{\frac{2}{3}}} \times \\ &\times \pi_{\Lambda}^{(\epsilon_2)}(p_1) \frac{|\widehat{\phi}_2(p_1-p_2)|}{E(p_1)^{\frac{1}{6}}E(p_2)^{\frac{1}{2}}} \pi_{\Lambda}^{(\epsilon_3)}(p_2) \frac{|\widehat{\phi}_3(p_2-p_3)|}{E(p_2)^{\frac{1}{2}}E(p_3)^{\frac{1}{2}}} \pi_{\Lambda}^{(\epsilon_4)}(p_3) \frac{|\widehat{\zeta}(p_3-p)|}{E(p_3)^{\frac{1}{2}}E(p)^{\frac{1}{6}}} \pi_{\Lambda}^{(\epsilon_1)}(p), \end{aligned}$$

so that, using (A.7) as above,

$$\begin{aligned} \int_{\mathbb{R}^3} |\widehat{Q_{3,\Lambda}^{\epsilon,\delta}\zeta}(p, p)| dp &\leq (K \log 3)^m \|M_{\phi_1}\|_{\mathfrak{S}_2} \left\| \frac{\pi_{\Lambda}^{(\epsilon_2)}}{E^{\frac{1}{6}}}(-i\nabla)\psi_2(x) \frac{\pi_{\Lambda}^{(\epsilon_3)}}{E^{\frac{1}{2}}}(-i\nabla) \right\|_{\mathfrak{S}_6} \\ &\times \left\| \frac{\pi_{\Lambda}^{(\epsilon_3)}}{E^{\frac{1}{2}}}(-i\nabla)\psi_3(x) \frac{\pi_{\Lambda}^{(\epsilon_4)}}{E^{\frac{1}{2}}}(-i\nabla) \right\|_{\mathfrak{S}_6} \left\| \frac{\pi_{\Lambda}^{(\epsilon_4)}}{E^{\frac{1}{2}}}(-i\nabla)\zeta(x) \frac{\pi_{\Lambda}^{(\epsilon_1)}}{E^{\frac{1}{6}}}(-i\nabla) \right\|_{\mathfrak{S}_6}, \end{aligned} \quad (\text{C.14})$$

where

$$\widehat{M}_{\phi_1}(p, q) = \frac{|\widehat{\phi}_1(p-q)|}{E(p+q)^{\frac{2}{3}}} |P_+^0(p)P_-^0(q)|.$$

The operator M_{ϕ_1} was estimated in Lemma 14 of [15] by $\|M_{\phi_1}\|_{\mathfrak{S}_2} \leq A\|\nabla\phi_1\|_{L^2}$, where A is some universal constant. By the definition of ϕ_1 , we obtain

$$\|M_{\phi_1}\|_{\mathfrak{S}_2} \leq 4\pi A\|(1+\mathcal{U})^m\mu_1\|_{\mathcal{C}}. \quad (\text{C.15})$$

As for the other terms in the right-hand side of (C.14), we argue as before and, by (C.14) and (C.15), we obtain

$$\int_{\mathbb{R}^3} |\widehat{Q_{3,\Lambda}^{\epsilon,\delta}\zeta}(p, p)| dp \leq \frac{A(K \log 3)^m}{\Lambda^{\frac{n(\epsilon)}{24}}} \|(1+\mathcal{U})^{-m}\zeta\|_{\mathcal{C}'} \prod_{j=1}^3 \|(1+\mathcal{U})^m\mu_j\|_{\mathcal{C}}, \quad (\text{C.16})$$

where A denotes some universal constant. All the terms in the right-hand side of (C.13) are similar to the one corresponding to $\delta = (1, -1, -1, -1)$. In particular, (C.16) holds when $Q_{3,\Lambda}^{\epsilon,\delta}\zeta$ is replaced by $Q_{3,\Lambda}^{\epsilon}\zeta$. By duality, this completes the proofs of (C.12) and of Step 3. This ends the proof of Proposition 3.2.

References

- [1] J. Avron, R. Seiler, and Barry Simon. The index of a pair of projections. *J. Funct. Anal.*, 120(1):220–237, 1994.
- [2] J.D. Bjorken and S.D. Drell. *Relativistic quantum fields*. McGraw-Hill, New York-Toronto-London-Sydney, 1965.
- [3] É. Cancès and M. Lewin. The dielectric permittivity of crystals in the reduced Hartree-Fock approximation. *Arch. Ration. Mech. Anal.*, 197:139–177, 2010.
- [4] P. Chaix and D. Iracane. From quantum electrodynamics to mean field theory: I. The Bogoliubov-Dirac-Fock formalism. *J. Phys. B*, 22:3791–3814, 1989.
- [5] P.A.M. Dirac. The quantum theory of the electron. II. *Proc. Roy. Soc. A*, 118:351–361, 1928.

- [6] P.A.M. Dirac. A theory of electrons and protons. *Proc. Roy. Soc. A*, 126:360–365, 1930.
- [7] P.A.M. Dirac. Theory of electrons and positrons. 1933. Nobel lecture delivered at Stockholm.
- [8] P.A.M. Dirac. Théorie du positron. volume XXV of *Solvay Report*, pages 203–212. Gauthier-Villars, Paris, 1934.
- [9] F.J. Dyson. The S matrix in quantum electrodynamics. *Phys. Rev. (2)*, 75:1736–1755, 1949.
- [10] F.J. Dyson. Divergence of perturbation theory in quantum electrodynamics. *Phys. Rev.*, 85(4):631–632, 1952.
- [11] E. Engel. Relativistic density functional theory: foundations and basic formalism. In P. Schwerdtfeger, editor, *Relativistic electronic structure theory, part 1. Fundamentals*, pages 524–624. Elsevier, Amsterdam, 2002.
- [12] E. Engel and R. M. Dreizler. Field-theoretical approach to a relativistic Thomas-Fermi-Dirac-Weizsäcker model. *Phys. Rev. A*, 35:3607–3618, 1987.
- [13] P. Gravejat, M. Lewin, and É. Séré. Ground state and charge renormalization in a nonlinear model of relativistic atoms. *Commun. Math. Phys.*, 286(1):179–215, 2009.
- [14] W. Greiner, B. Müller, and J. Rafelski. *Quantum electrodynamics of strong fields*. Texts and Monographs in Physics. Springer-Verlag, Berlin, first edition, 1985.
- [15] C. Hainzl, M. Lewin, and É. Séré. Existence of a stable polarized vacuum in the Bogoliubov-Dirac-Fock approximation. *Commun. Math. Phys.*, 257(3):515–562, 2005.
- [16] C. Hainzl, M. Lewin, and É. Séré. Self-consistent solution for the polarized vacuum in a no-photon QED model. *J. Phys. A, Math. Gen.*, 38(20):4483–4499, 2005.
- [17] C. Hainzl, M. Lewin, and É. Séré. Existence of atoms and molecules in the mean-field approximation of no-photon quantum electrodynamics. *Arch. Rat. Mech. Anal.*, 192(3):453–499, 2009.
- [18] C. Hainzl, M. Lewin, É. Séré, and J.-P. Solovej. A minimization method for relativistic electrons in a mean-field approximation of quantum electrodynamics. *Phys. Rev. A*, 76:052104, 2007.
- [19] C. Hainzl, M. Lewin, and J.-P. Solovej. The mean-field approximation in quantum electrodynamics. The no-photon case. *Commun. Pure Appl. Math.*, 60(4):546–596, 2007.
- [20] C. Hainzl and H. Siedentop. Non-perturbative mass and charge renormalization in relativistic no-photon quantum electrodynamics. *Commun. Math. Phys.*, 243(2):241–260, 2003.
- [21] C. Itzykson and J.-B. Zuber. *Quantum field theory*. Monographs and Textbooks on Pure and Applied Physics. McGraw Hill, New-York, 1985.
- [22] L.D. Landau. *On the quantum theory of fields. Bohr Volume*. Pergamon Press, Oxford, 1955. Reprinted in *Collected papers of L.D. Landau*. Pergamon Press, Oxford, 1965.
- [23] L.D. Landau and I. Pomerancuk. On point interaction in quantum electrodynamics. *Dokl. Akad. Nauk SSSR (N.S.)*, 102:489–492, 1955.

- [24] E. H. Lieb and H. Siedentop. Renormalization of the regularized relativistic electron-positron field. *Commun. Math. Phys.*, 213(3):673–683, 2000.
- [25] W. Pauli and M. E. Rose. Remarks on the polarization effects in the positron theory. *Phys. Rev. II*, 49:462–465, 1936.
- [26] P.-G. Reinhard, W. Greiner, and H. Arenhövel. Electrons in strong external fields. *Nuclear Physics A*, 166:173–197, 1971.
- [27] E. Seiler and B. Simon. Bounds in the Yukawa₂ quantum field theory: upper bound on the pressure, hamiltonian bound and linear lower bound. *Commun. Math. Phys.*, 45:99–114, 1975.
- [28] R. Serber. Linear modifications in the Maxwell field equations. *Phys. Rev. (2)*, 48:49–54, 1935.
- [29] D. Shale and W.F. Stinespring. Spinor representations of infinite orthogonal groups. *J. Math. Mech.*, 14:315–322, 1965.
- [30] B. Simon. *Trace ideals and their applications*, volume 35 of *London Mathematical Society Lecture Notes Series*. Cambridge University Press, 1979.
- [31] J.-P. Solovej. Proof of the ionization conjecture in a reduced Hartree-Fock model. *Invent. Math.*, 104(2):291–311, 1991.
- [32] B. Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.
- [33] E.A. Uehling. Polarization effects in the positron theory. *Phys. Rev. (2)*, 48:55–63, 1935.